

NEW STRAINS "fil", STRESS-RESISTANT UNDER FERMENTATION
AND/OR GROWTH CONDITIONS

5 The present invention concerns new eukaryotic strains, preferably yeast strains, having the new fil phenotype, i.e. having the unexpected property of conserving or keeping good stress resistance in fermentation and/or growth phase, while conserving normal respiratory and fermentation metabolism on fermentable sugars such as glucose.

The invention also concerns a process for obtaining such strains.

10 The invention also concerns the use of such strains for obtaining baker's yeast with higher resistance to drying, better adapted for the preparation of frozen dough and/or for other uses where good stress resistance during the fermentation phase is required.

15 Yeasts of the Saccharomyces genus are used as fermentation agents in baking, brewing, winemaking, distillery and other fields. Their industrial use is based on their ability to produce carbon dioxide from sugars such as glucose, fructose, sucrose or maltose, present or added in the dough or in the wort. Fermentation ability is an important criterion of quality for yeast.

20 The selection of strains, the manufacturing conditions of living or active yeast have been optimized during the years so as to obtain yeast having good fermentation ability and good stress resistance under certain conditions. Unfortunately, although yeast cells harvested under conditions equivalent to those of a stationary phase have a high level of resistance to different types of stress such as heat, freezing and drying, this stress resistance is lost when a fermentation phase is initiated by
25 adding fermentable sugars. The cells then rapidly lose their stress resistance property which causes a reduction in their fermenting power under stress conditions, this being a major disadvantage in the majority of the industrial uses for yeast.

The present invention concerns new eukaryotic strains and particularly new yeast strains obtainable by mutation (= mutagenesis), or by transformation with
30 recombinant DNA, called "fil" strains. The use of lower-case letters indicates that they do not have the FIL phenotype which is the normal phenotype, and stands for

“ Fermentation-Induced Loss of stress resistance ” which means loss of stress resistance induced by the fermentation. The new fil strains have a phenotype “ deficient in Fermentation-Induced Loss of stress resistance ” i.e. they are deficient in the loss of stress resistance which is induced by fermentation, without loss or significant alteration of their gassing power or fermentation activity and/or growth performance. In other words, the new strains conserve or keep in active metabolism phase a high level of stress resistance, comparable to that of cells which are not in fermentation or growth, i.e. are in stationary phase. This new property of the new strains is all the more unexpected insofar as the maintenance of good stress resistance is obtained without simultaneous significant loss of their growth and fermentation abilities.

All unicellular eukaryotic organisms (yeast, molds ...) are confronted with stress conditions whereas their industrial use requires them to be in active metabolic phase or to be able to rapidly achieve a (very) active metabolism. These two requirements, stress-resistance and active metabolism, have always been considered as contradictory and contrary to the natural biological equilibrium. For example, Attfield, in his review concerning the stress-resistance of yeasts of the *Saccharomyces* genus, published in December 1997 in “Nature Biotechnology”, 15, pp.1351-1357, writes that the reconciliation of these two requirements is contrary to biological design, i.e. to fundamental natural equilibrium or still to the original biological concept or definition of yeast strains. Consequently, the obtaining of new eukaryotic strains, particularly yeasts, having the unexpected properties of the fil phenotype, represents significant progress. If the example of baker's yeast is taken, it is well known that the more a cellular biomass is harvested under conditions close to exponential growth phase, the higher will be its fermenting power, but the lower will be its resistance to stress from drying or freezing. It is also well known that even if baking dough destined for freezing is seeded with a biomass harvested under conditions close to stationary phase, and consequently resistant to stress, the renewal or restarting of fermentation in the dough right from kneading and during the entire period before complete core freezing of the dough will induce a loss of this stress resistance. The new strains having the fil phenotype allow progress in the resolving of the difficulties inherent in the fundamental biological equilibrium of eukaryotic cells.

A lot of significant work has been performed in recent years to understand the mechanisms of action of stress factors on the cell and the nature of the cellular response, particularly in the case of *Saccharomyces cerevisiae*, when the yeast is exposed to different chemical or physical stresses. In general, the resistance appears to be a complex phenomenon, implying numerous factors which play distinctive roles. These factors, their mechanisms of action and their importance in relation to one another are still not well understood.

It is well known that cells of the yeast *Saccharomyces cerevisiae* become resistant to stress during stationary phase or when they are cultured or grown at a low growth rate on a non-fermentable carbon source or with a limited supply of fermentable sugar and/or nitrogen. However, this stress resistance disappears when a fermentable sugar such as glucose or maltose is supplied to the cells, which then enter rapid fermentation or active growth phase (Attfield, 1997, Nature Biotechnology, 15, pp.1351-1357 ; de Winde et al., 1997, Yeast Stress Responses, Ed. Springer, pp.7-52 ; Werner-Washburne et al., 1993, Microbiol.Rev., 57, pp.383-401).

The Ras-cAMP (Ras-cyclic Adenosine monophosphate proteins) metabolic pathway is known for its dramatic influence or role on the resistance to different types of stress in yeast cells. This has been demonstrated with respect to heat resistance (Iida, 1988, Mol. Cell. Biol., 8, pp.5555-5560 ; Matsumoto et al., 1985, Yeast, 1, pp.15-24 ; Shin et al., 1987, Mol. Cell. Biol., 7, pp.244-250), resistance to successive freezing and thawing steps (Park et al., 1997, Appl. Envir. Microbiol., 63, pp.3818-3824), and with respect to resistance to salt (Hitara et al., 1995, Mol. Gen. Genet., 249, pp.257-264).

The level of cAMP, and consequently that of protein kinase A activity in yeast cells is controlled by this elaborate and complex pathway (Broach and Deschenes, 1990, Adv. Cancer Res., 54, pp.79-139 ; Thevelein, 1991, Mol. Microbiol., 5, pp.1301-1307 ; Thevelein, 1992, Antonie Leeuwenhoek, J. Microbiology, 62, pp.109-130). In yeast, cAMP is synthesized by an enzyme, adenylate cyclase, which is encoded by the *CYR1/CDC35* gene (Kataoka et al., 1985, Cell, 43, pp.493-505). The level of cAMP is, in particular, regulated and hydrolyzed by two phosphodiesterases, encoded by the genes *PDE1* and *PDE2* (Nikawa et al., 1997, Mol. Cell. Biol., 7, pp.3629-3636 ; Sass et al., 1986, Proc. Natl. Acad. Sci. USA,

83, pp.9303-9307). Furthermore, the activity of adenylate cyclase is strongly dependent on the activity of the Ras proteins (Toda et al., 1985, Cell, 40, pp.27-36).

The Ras proteins are G proteins. They are active when bound or linked to a GTP (Guanosine TriPhosphate), and inactive when bound to a GDP (Guanosine DiPhosphate). The exchange of GDP with GTP on Ras proteins is stimulated by the guanine nucleotide exchange proteins Cdc 25 and Sdc 25 (Boy-Marcotte et al., 1996, Mol. Biol. Cell, 7, pp.529-539 ; Camonis et al., 1986, EMBO J., 5, pp.375-380). The Ras proteins have an intrinsic GTPase activity which is stimulated by the proteins Ira1 and Ira2 and which is responsible for the downstream regulation of their activity. cAMP activates cAMP-dependent protein kinase A (protein kinase A activated by cAMP hereafter referred to as PKA), which is composed of three catalytic subunits encoded by the genes *TPK1*, *TPK2* and *TPK3* and of a regulatory subunit encoded by the gene *BCY1* (Toda et al., 1987, Mol. Cell. Biol., 7, pp.1371-1327 ; Toda et al., 1987, Cell, 50, pp.277-287). The interaction between cAMP and the inhibiting subunit Bcy1 releases Bcy1 from the complex with the said catalytic subunits, thus activating them. These activated catalytic subunits phosphorylate a certain number of target proteins of which some have been identified, such as trehalase. The activity of PKA (protein kinase A activated by cAMP) is essential for the growth of yeast cells. When PKA activity is, in one manner or another, greatly reduced, e.g. by a severe reduction of cAMP level, the cells stop their growth and permanently enter stationary phase.

In other words, PKA, when it is activated, leads to growth of the yeast and is a mediator of different metabolic regulation processes, leading notably to a rapid decrease in trehalose content and a rapid decrease in heat shock proteins content, i.e. to disappearance of factors which favour stress resistance. On the contrary, when PKA activity is greatly reduced, the cells enter stationary phase and acquire high stress resistance. As indicated above, this has been demonstrated for different types of stress.

It is the study of these mechanisms of the complex Ras-cAMP metabolic pathway which has led Attfield review to establish that the obtaining of a phenotype corresponding to the maintenance of high stress resistance for cells in active metabolism after inoculation onto a medium containing fermentable sugars would be contrary to the “ biological design ” of the strains, i.e. against the natural equilibrium.

Mutants in the Ras-cAMP-protein kinase A pathway have been identified, constitutively having a high stress resistance during growth. This has been shown concerning heat resistance (Cameron et al., 1988, Cell, 53, pp.555-566 ; Hottiger et al., 1989, FEBS Lett., 255, pp.431-434 ; Shin et al., 1987, Mol. Cell. Biol., 7, pp.244-250) and freezing/thawing resistance (Park et al., 1997, Appl. Envir. Microbiol., 63, pp.3818-3824). However, these mutants have a much longer latent or lag phase at the start of fermentation and a reduced growth rate (Ma et al., 1997, Microbiol., 143, pp.3451-3459; Iida, 1988, Mol. Cell Biol., 8, page 5559) , which excludes their use in industry, particularly in bakers' yeast where a rapid onset of fermentation is essential.

Certain of these mutants have other properties important for use in industry affected in a negative manner. For example, *ras2* mutants are incapable of using a nonfermentable carbon source for their growth, ethanol for example (Tatchell et al., 1985, Proc. Natl. Acad. Sci., 82, pp.3785-3789 J.F. Canon and al., Genetics, 113, pp.247-264, June 1986). It is excluded to use a yeast having a deletion of the gene *RAS2* as a baker's or bread-making yeast strain because the assimilation of ethanol is necessary for the growth of baker's yeasts. In other words, it seems that these mutants constitutively conserve a high stress resistance since they are incapable of entering a truly active metabolic phase.

On the contrary, deregulated mutants having a high level of cAMP or a non limited PKA activity exhibit a very weak level of both heat shock proteins and trehalose, irrespective of the culture conditions and thus also in stationary phase.

The study of these mutants which have no industrial significance confirms the conclusions of the general review by Attfield already cited above concerning stress resistance, according to which it appears unlikely to obtain industrially useful strains by classical genetics and it is thus necessary to turn to recombinant DNA technologies. However, the theoretical data allowing such an approach are insufficient for obtaining the desired new result. This is particularly illustrated by the high amount of work which was performed in the past on deletion of the gene(s) coding for a trehalase.

It is well known that in bakers' yeast a high stress resistance, e.g. to heat, to freezing or to high pressures, is correlated to an elevated trehalose content (Attfield, 1997, Nature Biotechnology, 15, pp.1351-1357 ; De Virgilio et al, 1994, Eur. J.

Biochem., 219, pp.179-186 ; Iwakashi et al., 1997, Lett. Appl. Microbiol., 25, pp.43-47 ; Wiemken, 1990, Antonie Leeuwenhoek, J. Microbiology, 58, pp.209-217). Trehalose is a disaccharide present in high concentrations in a number of living organisms in nature (Van Laere, 1989, FEMS Microbiol. Rev., 63, pp.201-210 ; Wiemken 1990, Antonie Leeuwenhoek, J. Microbiology, 58, pp.209-217). It possesses remarkable and apparently specific properties of protection against aggressive treatments for a whole series of biological structures (Crowe et al., 1992, Anhydrobiosis Annu. Rev. Physiol., 54, pp.579-599). Trehalose is rapidly accumulated during the phase preceding the death of yeast cells.

Initiation of fermentation by the addition of a fermentable carbon source is associated with a rapid mobilization of trehalose (van der Plaats, 1974, Biochem. Biophys. Res. Commun., 56, pp.580-587), i.e. with its metabolic degradation and its rapid disappearance. Thus, a logical approach for maintaining stress resistance during the start of fermentation has been to clone and delete the *NTH1* gene, encoding for neutral trehalase (Kopp et al., 1993, J. Biol. Chem., 268, pp.4766-4774) which is the enzyme responsible for the mobilization of trehalose, so as to maintain the high trehalose level of the cells. It has been claimed that the stress resistance of yeast could be improved by the deletion of this gene (patents or patent applications EP 0451896 - Hino et al., EP 0838520). However, by impeding or preventing the mobilization of trehalose by the deletion of the *NTH1* gene, the rapid loss of stress resistance during the start of fermentation is not avoided (Van Dijck et al., 1995, Appl. Environ. Microbiol., 61, pp.109-115). This is also true concerning the non expression of the gene *ATH1* claimed in the patent application WO 97/01626. The simple deletion of one or all the genes coding for a trehalase is not by itself capable to solve the problem subject of the present invention.

This is probably due, in particular, to the action of other resistance factors such as heat shock proteins which disappear so rapidly at the start of fermentation (Crauwels et al., 1997, Microbiol. 143, pp.2627-2637 ; de Winder et al., 1997, Yeast Stress Responses, Ed. Springer, pp.7-52 ; Praekelt et Maecock, 1990, Mol. Gen. Genet., 223, pp.97-106 ; Werner-Washburne et al., 1989, J. Bacteriol., 171, pp.2680-2688).

Thus, the modification of the metabolism of trehalose by genetic engineering methods has not allowed the improvement of the stress resistance of yeast and has not given practical results for the development of industrial strains resistant to stress during fermentation phase (Attfield, 1997, Nature Biotechnology, 15, pp.1351-1357).

Another approach using recombinant DNA or genetic engineering techniques has been to try to increase the stress resistance of yeast by production of antifreeze proteins present in the blood of certain fishes inhabiting very cold waters. McKown et al., (Cryobiology, 1991, 28, pp.474-482) have expressed the gene encoding for an antifreeze protein in *Saccharomyces cerevisiae* so as to make it produce an intracellular chimeric antifreeze protein. However, this approach has not given satisfactory results since the survival rate of the yeast after freezing is still very low.

A large number of Japanese research activities can also be cited (patent applications or patents EP 0196233 - US4,547,374 - EP 0388262) dated of the 1980-1989 decade, consisting in selecting unconventional strains for bread-making fermentation but having interesting freeze resistance properties, and crossing them with baker's yeast. This approach has yielded limited results not solving the problem subject of the invention and none of these strains was used to produce baker's yeast commercialized on the European or American market in 1998.

Thus it has not been possible until now to obtain yeast having the property of conserving a high stress resistance simultaneously with good growth and good fermenting activity.

In conclusion it can be said, in agreement particularly with the conclusions of the general review on yeast stress published in Nature Biotechnology in 1997, that :

the major problem is that the natural response of the cells, in the presence of a fermentable substrate, is to pass into active metabolic phase and thus to rapidly decrease their stress resistance factors, whereas industrial conditions necessitate active metabolism and high stress resistance ;

this problem was not resolved;

classic genetics have only contributed a limited improvement of the

resistance of cells in industrial use conditions ;

a solution of this unsolved problem could only be expected via genetic engineering technologies.

Unfortunately, despite the complete knowledge of the yeast genome which has been entirely sequenced, and the knowledge which has accumulated regarding the functions of the genes but which is still very incomplete, it is only partially known how the industrially crucial properties are genetically and physiologically governed. The present state of the art is thus insufficient to allow adequate genetic manipulation to arrive at a solution of the major problem defined above. This is all the more true insofar as the regulatory pathways concerned, such as the Ras-cAMP pathway are very complex as illustrated above, and are probably numerous. The Ras-cAMP-PKA pathway is not the only metabolic (regulatory) pathway to be considered.

The present invention, in a surprising manner, has resolved these problems in a simple and efficient manner. It demonstrates that it is possible, contrary to that which was generally admitted, to obtain strains having a phenotype which was believed not to be capable of existing. This phenotype contrary to "biological design" (i.e. to natural equilibrium) has been called the fil phenotype.

BRIEF DESCRIPTION OF THE INVENTION

The present invention concerns a process for obtaining new eukaryotic strains, preferably new yeast strains, and even more preferentially strains of *Saccharomyces cerevisiae* conserving stress resistance in the presence of fermentable sugars such as glucose, characterized by the fact that it comprises the following steps :

a classic mutagenic treatment is carried out on the cells of a starting or original strain,

the cells having undergone the said mutagenic treatment are cultured or grown until they reach the stationary phase,

the said cells in stationary phase are incubated in the presence of at least one fermentable sugar selected from the group comprising glucose, maltose, and sucrose,

this sugar being present in a quantity such that the cells enter an active metabolic state (fermentation and/or growth) of this sugar,

said cells in active metabolic state are subjected to one or several stresses leading to a mortality rate of at least 99% with respect to the starting population,

5 the surviving cells are isolated and

those of the surviving cells which respond to the following criteria which characterize the fil phenotype are selected, i.e. :

- a growth, evaluated by biomass production or production yield of biomass over sugar consumed in a given time or by a maximal growth rate, under identical culture conditions, at least equal to 80% of the starting or control strain, and preferably at least equal to 90% of the starting or control strain,
- a CO₂ release, or a metabolite production, in identical conditions, at least equal to 80%, and preferably at least equal to 90% of the starting or control strain,
- a stress resistance, corresponding to a survival rate at least 2 times higher, preferably at least 3 times higher, more preferentially at least 5 times higher, and even more preferentially at least 10 times higher than the survival rate of the starting strain, under identical phase conditions corresponding to a growth or active metabolism followed by a heat shock of at least 20 minutes at 52°C, and/or at least 1.5 times higher, preferably at least 2 times higher, more preferably at least 3 times higher, and even more preferentially at least 5 times higher than the survival rate of the starting strain, under identical conditions of growth or fermentation phase followed by freezing for a period of at least 24 hours at -20°C or at a lower temperature,
- maintenance of these properties after repeated cultures or cultivations on non selective medium, such as YPD medium, so as to verify that the fil phenotype obtained by the mutation is perfectly stable and permanent.

Preferably, the absence of all hampering properties possibly accompanying the phenotype fil will be checked. For instance, these hampering properties can be the formation of inopportune secondary metabolites, or the loss of the capacity of assimilation or fermentation of certain compounds or substrates.

30 According to a particular aspect of the invention, the selected fil strains preferably have the property of conserving, in growth and/or fermentation phase on fermentable sugars, at least 50%, preferably at least 60%, more preferentially at least

70%, and even more preferentially at least 80% of their survival rate with respect to the survival rate in stationary phase measured under the same conditions after a heat or freeze shock.

In a particular embodiment of the process for obtaining the new fil strains,
 5 the cells obtained after mutagenesis treatment and in stationary phase are introduced into pieces of dough of 0.5 g consisting of water (about 42.5%), flour (about 56.5%), NaCl (about 1%), at a level of $4 \cdot 10^8$ cells per g of dough. Said pieces of dough are subjected to a first fermentation of 30 minutes at 30°C, then are subjected to at least 100 cycles of freezing/thawing.

10 Preferably, the process of obtaining the new eukaryotic strains, subject of the invention, is applied directly to industrial strains. The selection tests used correspond to the stress encountered by the said eukaryotic strain in the process(es) of production used and to the characteristics of performance of the said strain. The strain having the selected fil phenotype is a strain having the characteristics justifying its use
 15 in the industrial or craft (= artisan) production coupled with a better resistance to the encountered stresses. An industrial strain is a strain really used in optimized and competitive industrial production. If the starting (= original) strain is a baker's yeast strain, this strain is then a strain actually used by a specialized yeast producer for the selling on the market of baker's yeasts, or a strain having equivalent properties.

20 A laboratory strain is a model strain allowing a good understanding of the studied phenomena, but which does not have all the properties necessary to industrial strains. A yeast model strain is a true haploid or a true diploid of *Saccharomyces cerevisiae*, which can contain a certain number of auxotrophic markers. The industrial yeast strains are usually polyploids.

25 In a particular embodiment of the process for obtaining new fil strains, wherein an industrial fil mutant strain carrying several mutations, then one can improve the said mutant in the following manner: Process according to claim 3, wherein an industrial fil mutant carrying several mutations is obtained and wherein :

- the segregants issued from this industrial mutant are crossed with a laboratory
 30 haploid strain to select the segregant issued from this industrial mutant giving to the polyploids obtained with the laboratory strain an improvement in the required properties;

- the segregants thus selected are crossed one with the other;
- the polyploids obtained are selected according to the above-defined criteria of fil phenotypes.

The present invention also concerns new eukaryotic strains, preferably new
 5 yeast strains belonging to the *Saccharomyces* genus, preferably *Saccharomyces cerevisiae*, having the fil phenotype, obtainable by the process described above or one of its embodiments.

The present invention concerns:

on the one hand new laboratory yeast strains, new haploids or segregants or
 10 segregeants of laboratory strains or industrial strains having the fil phenotype, these new strains being essentially tool or model strains for the construct of industrial strains possessing the fil phenotype

on the other hand, and this is the main subject of the invention new
 15 industrial yeast strains, preferably new industrial baker's yeast strains possessing the fil phenotype.

An advantageous particularity of the present invention is to obtain
 industrial eukaryotic fil strains not GMO, that is not Genetically Modified
 microOrganisms in the meaning of the directive CEE 90/220 for example. Considering
 the unjustified European reluctance, in the light of the precise regulation existing in
 20 Europe towards GMOs, it is significant that the process subject of the invention allows to construct new industrial baker's yeast strains that possess the fil phenotype and that are not genetically modified, i.e. that are non-GMO.

In particular, these new yeast strains presenting the phenotype fil preferably
 have a survival rate, in growth phase on fermentable sugar, of at least 50%, preferably
 25 at least 60%, more preferably at least 70%, and even more preferably at least 75%, after heat treatment of 20 minutes at 52°C, the growth phase being defined :

- for laboratory strains (true haploid or diploid strains, generally auxotrophic) and all segregants of industrial strains, as a reculturing on fermentable sugar (glucose) for 30 minutes at 30°C after stationary phase, i.e. as a cultivation of
 30 stationary cells for 30 minutes at 30°C;
- for industrial strains (aneuploid and polyploid strains), as a cultivation

of stationary cells on fermentable sugar (glucose) for 10 minutes at 30°C after stationary phase.

The new yeast strains according to the invention are all new yeast strains of the laboratory type (true haploid or diploid strains, generally auxotrophic) and all
 5 segregants of industrial strains, whose stability to freezing in pieces of dough containing 20 g of flour, 15 g of water, 1 g of sucrose, 0.405 g of NaCl, 0.06 g of $(\text{NH}_4)_2\text{SO}_4$ and 160 mg of dry matter of the considered strain is at least higher than 60%, preferably at least higher than 70% and more preferably at least higher than 80%, stability being defined by the ratio between the release of CO_2 at 30°C after 1 month
 10 (30 days) of conservation at -20°C and the release of CO_2 at 30°C after 1 day of conservation at -20°C.

The new yeast strains according to the invention are also all new industrial yeast strains (aneuploid or polyploid strains), whose stability to freezing in pieces of dough containing 20 g of flour, 15 g of water, 0.405 g of NaCl, 0.06 g of $(\text{NH}_4)_2\text{SO}_4$
 15 and 160 mg of dry matter of the considered strain, measured by the ratio between the release of CO_2 at 30°C after 1 month (30 days) of conservation at -20°C and the release of CO_2 at 30°C after 1 day of conservation at -20°C, is at least higher than 80%, preferably at least higher than 85% and more preferably at least higher than 90%.

These freeze-stability tests correspond respectively to tests C2 and C1
 20 described hereafter. Before freezing at -20°C, these pieces of dough are incubated at 30°C for 30 minutes (test C1 on industrial yeast) or 60 minutes (test C2 on laboratory strains or segregants).

Preferably, the new yeast strains according to the invention allow the obtaining of dry yeast from a biomass harvested in exponential growth phase or in
 25 phase close to exponential growth phase, having a loss or decrease of released gas (= gassing power) after drying at most equal to 67%, preferably at most equal to 50% of the loss of released gas after drying of yeasts obtained using the corresponding starting or original strain (non mutated) or a control strain having the same characteristics.

30 The present invention also concerns the new strains having the fil phenotype :

- PVD1150 = M5 *fil1* deposited at C.N.C.M. 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-2031 (contaminated strain) and I-2203, in accordance with the Budapest Treaty.
- KL1 = W303 *fil2* deposited at C.N.C.M. 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-2032, in accordance with the Budapest Treaty.
- FD51 = HL816 *fil300* deposited at C.N.C.M. 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-2033, in accordance with the Budapest Treaty.
- FDH16-22 = HL822 *fil300* deposited at C.N.C.M. 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-2034, in accordance with the Budapest Treaty.
- AT25 = S47 *fil400* deposited at C.N.C.M. 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-2035, in accordance with the Budapest Treaty.
- AT28 = S47 *fil500* deposited at C.N.C.M. 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-2036, in accordance with the Budapest Treaty.

and all the new strains of the same kind, that is to say all the new strains having similar characteristics.

As the first deposit of the strain PVD1150 = M5 *fil1* revealed to be contaminated, it was deposited again on May 20th, 1999 under the number I-2203.

The invention also concerns a mutant or mutated gene or mutated genes which confer the phenotype *fil*, this or these genes being obtained by molecular biology techniques from eukaryotic *fil* strains obtained by the process of obtaining *fil* mutants. In particular, the said gene is the gene or genes which confer(s) the phenotype *fil* to one of the *fil* strains which exemplify the present invention and which have been deposited at the C.N.C.M. For example, the said gene is the gene *CDC35* = *CYR1* carrying a mutation conferring the *fil* phenotype.

Advantageously, the said mutation in the gene *CDC35* = *CYR1* is a change of a G base (guanine) into an A base (adenine) in the region of the gene *CDC35/CYR1* coding for the catalytic site of the enzyme, equivalent to a change of an acidic amino acid (glutamic acid) into a basic amino acid (lysine) at position 1682 of the protein. This mutation is responsible for the *fil* phenotype in the strain PVD1150 = M5 *fil1*.

This gene can also be the gene YDL 035C according to the nomenclature defined in the publication concerning the sequencing of the yeast genome, project published in Nature, 1992, 357, pp.38-44, hereafter called *GPR1*, carrying a mutation conferring the fil phenotype and more particularly the mutation of the strain KL1 =
 5 W303 *fil2*. This gene can also be the mutated gene or one of the mutated genes conferring the fil phenotype of one of the strains C.N.C.M. I-2033, I-2035, I-2036.

In a general manner, the present invention includes all genes encoding for a protein having similar or equivalent properties to the proteins encoded by the genes defined above as carrying a mutation conferring the fil phenotype, i.e. all genes
 10 carrying a mutation conferring the fil phenotype and belonging to the family of genes :

- encoding for a protein having a function comparable or equivalent to that of a protein encoded by one of the genes defined above as carrying a mutation conferring the fil phenotype in yeast or another eukaryote, a comparable function being defined as the commanding of the same mechanisms in the same metabolic pathway or an
 15 equivalent metabolic pathway,
- encoding for proteins associated with the protein encoded by one of the genes carrying a mutation conferring the fil phenotype,
- encoding for proteins having similar sequences, i.e. at least 60% homology or identity, preferably at least 70% homology and still more preferably at least 80%
 20 homology with the protein encoded by one of the genes carrying a mutation conferring the fil phenotype,

For example, this gene can encode for a protein associated with the protein encoded for by the gene *GPR1* defined above, and specifically, this gene can be the gene(s) *GPA2* of yeast carrying a mutation which confers the fil phenotype.

25 In general, the present invention is not limited to yeast strains and concerns all eukaryotic strains carrying a fil mutation i.e. having a fil phenotype.

Said eukaryotic strain is advantageously transformed in a manner so that at least certain of the alleles of a gene, capable once mutated of conferring the fil phenotype, carry the mutation conferring the said fil phenotype. Said eukaryotic strain
 30 is preferably an industrial strain and preferably an industrial yeast strain.

The invention also concerns the use of yeast strains mutated and selected for their fil phenotype obtainable by the process, subject of the invention, or the use of

strains genetically transformed in such a manner as to have the fil phenotype, for obtaining bread making yeast (= baker's yeast), in particular destined to the inoculation of frozen dough.

5 The present invention also concerns the use of the said mutated or transformed fil strains for obtaining dry bread making (= baker's) yeast.

The present invention also concerns the use of said mutated and selected or transformed fil strains for obtaining industrial brewer's yeast, winemaking yeast or yeasts destined for the production of alcohol.

10 In a general fashion, the present invention concerns the use of new eukaryotic strains having the fil phenotype in any industrial condition necessitating simultaneously properties of resistance to stress and of active metabolism of said eukaryotic strains.

DETAILED DESCRIPTION

15

The invention concerns new eukaryotic mutant strains called fil, preferably new fil yeast strains, which conserve or keep high stress resistance during an active growth or active fermentation phase on glucose and which have conserved the essential part of their metabolic properties (growth, production of primary or secondary metabolites). These characteristics of fil strains were considered as irreconcilable or incompatible and contrary to biological design. The search for such new strains consequently went against a preconceived opinion. The new process used for the obtaining of such eukaryotic strains having unexpected properties is based on the following steps :

20

- 25 • the cells are subjected to a known mutagenesis treatment or protocol;
 - the cells thus obtained are cultured or grown until they reach stationary phase, then are recultured or cultivated in the presence of a fermentable sugar such as glucose to be in active metabolism ;
 - said cells are subjected to a strong heat or freeze stress, i.e. a stress causing high lethality, so as to select those which have become resistant to stress in active metabolic phase ;
- 30

- the different successive steps described above or at least one of them may be repeated so as to obtain a survival rate with respect to the starting population equal to or lower than 1% and preferably to 1 per 1000, or even 1 for 1 million ;
- the surviving mutated or mutant strains (cells) thus obtained are tested to verify their resistance to heat or freezing in growth or fermentation phase, preferably they are also tested to check the non appearance of an undesired secondary property or the non disappearance of an interesting or useful secondary property, then are selected from among the resistant strains those strains which have substantially conserved their properties of growth, fermentation and/or synthesis of metabolites of industrial interest, preferably without appearance of hampering properties and without disappearance of useful properties ;
- the maintenance of properties corresponding to the fil phenotype is finally verified after cultivation of the said strain corresponding to a large number of generations under non selective conditions for cells.

This process of searching for fil strains may be applied to all eukaryotic strains or organisms, the tests optionally being adapted as a function of the characteristics of the said eukaryotic strain. It is noted that due to the fact that the resistances are generally crossed, the tests of resistance to heat and/or freezing are good selection tests for the search for any new strain resistant to a given stress that it will encounter during its industrial use. In order to simplify the language, these two tests are referred to below as tests of resistance to thermal shock.

In an entirely unexpected manner, since it was thought that such strains did not exist, the process thus described effectively led to the selection of several perfectly stable strains having these characteristics corresponding to the fil phenotype and notably to the selection of new industrial baker's (or bread-making) yeast strains, not GMO and directly usable to the commercialization of new baker's yeast strains.

The culture media used in the present invention are :

30	YPD medium	yeast extract	10 g/l
		bactopeptone	20 g/l
		glucose	20 g/l

	YPD-A medium	yeast extract	10 g/l
		bactopeptone	20 g/l
		glucose	20 g/l
		agar	20 g/l
5	YP medium	yeast extract	10 g/l
		bactopeptone	20 g/l
	SD-URA	Nitrogenous base medium free of aminoacid	
10		(Yeast Nitrogen Base DIFCO®)	6.7 g/l
		complementary mixture without uracil (CSM-URA, Bio 101®)	0.77 g/l
		glucose	20 g/l
		agar	15 g/l
15	R medium	glucose	1.5 g/l
		yeast extract	1.0 g/l
		MgSO ₄	0.7 g/l
		CaCl ₂	0.4 g/l
20		(NH ₄) ₂ SO ₄	2.0 g/l
		KH ₂ PO ₄	1.87 g/l
		K ₂ HPO ₄	1.1 g/l

The tests used in the present invention are :

25 TESTS FOR SCREENING OF MUTANTS RESISTANT TO THERMAL STRESS DURING ACTIVE FERMENTATION AND/OR GROWTH PHASE

Test T1 : Survival rate of cells after heat thermal shock

30 The cells are cultured or grown under stirring on YPD medium until stationary phase is obtained. These cells in stationary phase are washed, resuspended in ice-cold YP medium, so that the optical density at 600 nm is preferably comprised between 1 and 2 with respect to the medium. This suspension is then incubated at 30°C until said temperature is reached. Part of the suspension is then kept on ice in order to serve as a control for stationary phase cells. Glucose is added to the other part

35 to a final concentration of 100 mM. This addition is followed by an incubation of cells suspension at 30°C of 10 to 90 minutes and this other suspension part is then kept on ice at the end of the chosen time. Both suspensions are subjected, on the one hand, to a counting of the number of viable cells after suitable dilution and, on the other hand, to a thermal treatment of at least 20 minutes at 52°C followed by the same counting of the

number of viable cells. This counting is performed on YPD-A after two day incubation at 30°C.

A measurement of the stress resistance or survival rate on the one hand of cells in stationary phase and on the other hand of cells in active metabolism can thus be obtained. The survival rate is expressed as a ratio of the number of colonies formed in the thermally treated samples and the number of colonies formed in the control samples.

Test T2 : Survival rate of cells after cold thermal shock

Identical to Test T1 except for the stress applied : the suspension is incubated at -20°C to -30°C for 1 to 12 days.

Test T3 : Determination of the growth rate

The cells are cultured on YPD medium at 30°C under stirring at 180 rpm until stationary phase is obtained.

The growth, i.e. the proliferation of cells is monitored as a function of time by measurement of the absorbency (= optical density) at 600 nm with respect to non seeded medium.

It is expressed by the curve of development of the absorbency, i.e. by the change of optical density in function of time.

The growth rate is the rate of increase of the number of cells, i.e. the slope of the absorbency curve as a function of time.

Test T4 : Determination of growth rate in microplates

The cells are cultured on YP medium + glucose at 10 or 100 mM concentrations or on YP medium + beet molasses at 5 g/l at 30°C and in microtitration plates (microplates). 250 µl of medium are seeded in order to obtain 0.05 OD at 600 nm of cells in stationary phase. The microplates are shaken for 30 seconds of each minute and the absorbency at 600 nm is measured every 30 seconds. The growth rate corresponds to the slope of the absorbency curve, the μ_{\max} or maximum growth rate is then determined.

Dishes containing 100 grams of agar medium having the following composition:

are seeded or inoculated with the equivalent of 2 mg dry matter of yeast per dish. These dishes are incubated at 30°C for 20 to 40 hours. The final quantity of yeast dry matter produced in weight is measured in 10 to 20 dishes, generally 16 dishes.

15

To test the loss of the fermentative capacity, the glucose consumption of frozen cells is measured and compared the glucose consumption of the same non frozen cells but stored on ice. The residual fermenting power after freezing is then determined by the ratio between the glucose consumption of frozen cells and the glucose consumption of non frozen cells. This ratio corresponds to test T7, the glucose consumption of non frozen cells corresponds to test T6.

Afterwards, cells are resuspended in YP medium to obtain an Optical Density (OD) at 600 nm equal to 15. Samples of 0.03 ml are removed and frozen directly in a methanol bath of -25°C during 1 hour, then are stored at -30°C during 1 to 36 days then thawed at ambient temperature.

In parallel, samples are stored on ice and are used as unfrozen control cells. Thawed cells samples and control cells samples are 11 times diluted with YP-10mM glucose

and incubated at 30°C. 0.01 ml cell free samples are collected after 90 and 120 minutes. Residual glucose is measured by adding 0.2 ml Trinder Reagent® (supplier: Sigma). After 15 minutes of incubation at 30°C, the absorbance is measured at 505 nm. An almost linear correlation is found between the amount of active cells and the glucose consumption making it relatively simple to determine the residual fermentation capacity of the frozen samples.

TESTS FOR DETERMINING THE GASSING POWER OF YEAST

Tests A1, A20, A'1, A'20, C1 and C2, used for determining fermenting or gassing power of yeast, i.e. their capacity to produce CO₂, are performed with the help of a fermentometer from Burrows and Harrison, described in "Journal of the Institute of Brewing", 1959, LXV, 1, January-February, and are precisely defined in the following manner :

Test A1 : Fermenting power (fresh yeasts, industrial strains)

To 20 g of flour incubated at 30°C is added a weight of yeast corresponding to 160 mg dry matter, this yeast being suspended in 15 ml of water containing 27 g of NaCl per liter and 4 g of (NH₄)₂SO₄ per liter ; the suspension is mixed with a spatula for 40 seconds, so as to obtain a dough which is placed at 30°C ; thirteen minutes after the beginning of mixing, the recipient (pot) containing the dough is hermetically sealed ; the total quantity of carbon dioxide (CO₂) produced is measured after 60 and 120 minutes, this quantity is expressed in ml at 20°C and under 760 mm of mercury (Hg).

Test A20 : fermenting power (fresh yeasts, other strains)

Test identical to test A1 but the composition of the dough is modified as follows : 1 g (1 gram) of sucrose is added to the mixture of flour, water, yeast and salt, before carrying out the kneading. Furthermore, the release of CO₂ is measured after 120 and 240 minutes at 30°C (instead of 60 and 120 minutes for test A1).

Test A'1 : fermenting power (dry yeasts, industrial strains)

Test identical to test A1 but before mixing, the 160 mg of dry matter content of yeast, which has the form of dry yeast, is rehydrated for 15 minutes in

[illegible]

5

Test identical to test A'1, but 1 g of sucrose is added to the flour ; the total quantity of gas produced is measured over 240 minutes.

Test C1 : fermenting power (industrial strains after freezing)

Test identical to test A1 but it is necessary to prepare at least six pieces of dough per strain. The pieces of dough are prepared or made according to the conditions of test A1, but after kneading, the pieces are incubated at 30°C for 30 minutes. The prefermented pieces of dough are then immediately stored at -20°C and conserved at this temperature for periods from 1 day to 2 months. To measure the fermenting activity after conservation at -20°C, the frozen dough is placed in an incubator at 30°C ; after thirteen minutes, the pot containing the dough is hermetically sealed and the total quantity of CO₂ produced (expressed in ml at 20°C and under 760 mm Hg) is measured after 120 minutes with a fermentometer of Burrows and Harrison.

For a given strain, the reference release of CO₂ corresponds to the release of CO₂ of a dough sample stored at -20°C for a single day. The other pieces of dough are thawed at regular intervals (for example after 1 week, 2 weeks, 1 month (i.e. about 30 days), 1 ½ months and 2 months of storage at -20°C), and the release of CO₂ is measured so as to follow the evolution of the fermenting activity as a function of the duration of storage at -20°C.

25 The stability to freezing is defined as the ratio of CO₂ release during 2 hours after 1 month of freezing and the reference CO₂ release during 2 hours after 1 day of freezing.

Test C2 (other strains after freezing)

30 Test based on test C1 with the following modifications :

- 1 g of sucrose is added to the flour-water-salt-yeast mixture before kneading,

- the pieces of dough are prefermented 60 minutes at 30°C before freezing at -20°C,
- the CO₂ release is measured during 240 minutes at 30°C.

5 Test R (determination of ethanol assimilation)

100 ml of R medium are inoculated with stationary phase cells, previously grown on YP medium + glucose at 100 mM during 24 to 48 hours, so that the optical density (OD) of the inoculated culture is 0.05 OD at 600 nm. The inoculated R medium is then incubated at 30°C and stirred continuously at 180 rpm (revs per minute). The
 10 absorbency (optical density) at 600 nm is monitored with regular takings until obtaining a stable OD. Within these conditions, a strain, which does not assimilate ethanol reaches an OD about 4 times lower than a strain normally assimilating ethanol. Only the strains reaching an OD at least equal to 50% of the control strain (in general, the non mutated strain) are retained, and preferably at least 80%, and more preferably
 15 at least 90%.

All the tests described above for putting into practice of the invention are tests of a biological nature and their reproducibility from one laboratory to another often causes problems of a delicate nature. Consequently, they should usually be interpreted in a relative fashion with respect to a control. The tests should be
 20 conducted so as to reproduce the values indicated for the control, or preferably, the controls so as to have a scale of reproducible values. This conducting of the tests should be carefully and meticulously performed.

It is clear for the person skilled in the art that each step of the process should be adapted in accordance with the characteristics of the strains used and their
 25 lethality or survival rate in the different tests. The indications given hereafter concerning the carrying out of the different steps of the process, subject of the invention, are only examples for the application of the process to *Saccharomyces cerevisiae*.

The process object of the invention whose principle is given hereabove and
 30 which is intended to find new strains that simultaneously have an active metabolism and are stress resistant, may be carried out:

- on lab strains, i.e. on model strains whose genetic characteristics are well known and which generally contain auxotrophy markers;
- on industrial strains, in general much more complex on the genetic level, due to the fact that they are not true haploids or diploids, contrary to lab strains, said industrial strains having been selected on the basis of their industrial performances. Among the industrial strains, it is possible to distinguish:

- the industrial strains in their most stable state, in general in polyploid form,
- the segregants of industrial strains, i.e. the sexual forms which are in general less stable, but which can be used for the constructions carried out by classic genetics.

Within the frame of the research carried out in order to characterize the mutations providing the fil phenotype, the new process object of the invention will be preferably carried out on lab strains, preferably haploids.

The characterization of the gene or of the genes involved will be carried out to allow subsequent constructions of industrial strains, carrying the corresponding mutation or mutations.

Within the frame of research carried out in view of obtaining directly industrial strains, the process will be carried out directly either on segregants of industrial strains or directly on industrial strains. The use of segregants means that the fil segregants thus obtained will be subjected to constructions by classic genetics in order to recover all desired characteristic features of industrial strains, in addition to the fil phenotype.

The obtaining of fil industrial strains, not GMO, which means not Genetically Modified Organisms, is the preferred subject of the invention.

The main characteristics of these industrial strains are the following ones:

- preservation of the main properties of the industrial strains actually used because of their efficiency (performance) they are derived from
- significantly increased resistance to the stress conditions met in the processes of industrial or craft production they are used in
- contribution to a progress (like best productivity or lower industrial cost) or a best result linked to the fil phenotype.

According to its intrinsic definition, the fil phenotype cannot correspond for a yeast strain for example:

- to a deletion of *RAS1* or *RAS2* genes, considering the poor assimilation of ethanol such a deletion implies (J.F. Cannon and al., Genetics, 113, p.250, June 1986). The
5 fil strains must demonstrate that they are able to re-assimilate one part of the alcohol in the test R because this is a useful secondary property which must not disappear.
- to the *cyr1-T1* and *cyr2-T2* mutants described by Iida (Mol. Cell Biol., Dec. 1988, pp.5555-5560) considering the low growth rates of these mutants.
- to a particular sensitiveness of its multiplication or its fermentation to temperature.
- 10 • to a simple deletion of one or several genes encoding for one of the trehalases of yeast, the non synthesis or the low synthesis of trehalases in growth condition being an interesting factor as a secondary factor of the stress resistance, but insufficient by itself.
- to a transformation with DNA encoding for a protein with a SOD activity
15 (SuperOxide Dismutase) and with DNA encoding for a protein with catalase activity, because this corresponds to secondary factors with limited action, and to loss of efficiency or performance linked to the expression of these genes within the scope of the normal use of such yeasts.

A bread-making yeast strain, which after mutation, would give for instance a bad smell
20 to breads, a bad or abnormal taste to breads, due to a secondary metabolism affected by the mutation, would not be a strain corresponding to the fil phenotype, because according to its preferential definition given in the present invention, the fil phenotype corresponds to a yeast having not any secondary property undesirable for its use.

Advantageously, the mutagenic treatments used in the process according to
25 the invention are as follows. The cells of *Saccharomyces cerevisiae* yeast are cultured on YPD medium and are then subjected to a mutagenic treatment using a chemical agent such as EMS (Ethyl-Methyl-Sulfonate) or by ultraviolet light according to the classic protocols (Sherman et al., 1986, Cold Spring Harbor Lab Press ; Spencer et al., 1988, Yeast a practical approach, Ed. Campbell et Duffus). The conditions of the
30 mutagenesis are in general selected in order to obtain a survival rate of the cells of the order of 1 to 20%, preferably about 10%. The cells are then washed, suspended again in YPD medium and cultured until the stationary phase is obtained. Then a known

amount of the culture is taken, possibly washed, transferred on YPD medium and incubated at 30°C, for 30 to 90 minutes. According to a variant, it is possible to add to the culture in stationary phase 100 mM glucose and to incubate at 30°C for 30 to 90 minutes.

5 The culture is then subjected either to a heat shock by incubation between 52 and 65°C, preferably at 56°C for 30 minutes or more, or to a cold shock (freezing) by incubation between -20°C and -40°C for 1 to 3 days. The freezing induces only a small loss of viability of the yeast cells, the treatment is repeated up to 200 times, until a survival rate of the cells lower than 1 for 10000, preferably lower than 1 for 100000
10 and still more preferably lower than 1 for 1000000 is obtained.

In a particular embodiment of the selection protocol, the cells after mutagenesis are introduced in small pieces of dough of about 0.5 g in a proportion of 4.10^8 cells/g of dough, the latter being composed of water (42.5%), of flour (56.5%), of NaCl (1%) and subjected to fermentation for 30 minutes at 30°C in such a way that the
15 yeast cells leave the stationary phase, i.e. enter the fermentation phase. The pieces of dough are then successively frozen at -30°C and thawed at room temperature, up to 200 times in such a way that only a few hundred or a few thousand cells survive.

At this stage, the cells, which survived the thermic shock (heat or freezing) are mutants, which are stress resistant and of which it is necessary to check, on the one
20 hand, the persistence of the said property during fermentation or growth for several generations and, on the other hand, the maintenance of their growth ability and that of the production of metabolites.

In order to check the stress resistance in growth phase and/or in fermentation phase, the survival rates are measured according to the tests T1 and/or T2, and/or equivalent tests. The mutated strains are selected according to the increase of
25 their survival rate under an important stress during or after fermentation phase and/or growth phase like those according to tests T1 and T2 in comparison with the starting strain, and in that scope, from a general point of view, a survival rate of at least 1.5 times higher will be required. The mutated strains can also be selected by comparison
30 of their survival rate in growth phase and/or fermentation phase with respect to their survival rate in stationary phase, or with respect to the survival rate in stationary phase of the starting (original, non modified) strain.

Preferably, the selected mutated strains will present, after a growth and/or fermentation phase of the kind defined in tests T1 or T2 at least 50%, preferably at least 60%, more preferably at least 70%, and still more preferably at least 80% of their survival rate in stationary phase.

5 According to a variant, the resistance of the mutated strains which are obtained is measured, after freezing according to the tests C1 or C2 or according to equivalent tests, or still again the resistance against drying is measured using the ratio starting from the values obtained in tests A and A' with respect to the yeast before and after drying.

10 In order to check the conservation of the growth and fermentation properties, a comparison can be made between the starting strain and the mutated strain having a phenotype of stress resistance in metabolic active phase, in one of the tests T3, T4 or T5 and in one of the tests A or in similar tests which permit the measurement of the properties which are of interest for the considered strains. A conservation of at
15 least 80% of these properties is necessary to correspond to the fil phenotype.

Preferably, it will be checked that the mutation did not lead to any loss of an interesting secondary characteristic. For example, it will be checked by test R that the strain has a sufficient alcohol assimilation, at least 50% of that of the control strain. It will also be checked that there is no production of inopportune metabolite. For
20 example, it will be checked on the industrial baker's yeast strains that under the usual conditions of bread-making notably in Europe and in the USA, there is no bad taste or bad smell given to the breads.

Furthermore, it is necessary to check that the mutation is stable, in other words that the fil phenotype is stable. Consequently, all the above-defined verifications
25 should be repeated on the reisolated mutated strain after a great number of multiplications in non selective medium, as for example after 10 successive cultures of 2 days in YPD-A medium at 30°C or again after 10 cultures in YPD medium at 30°C, each culture being seeded by 1 per cent from the preceding culture.

The strains which have withstood with success these different selection
30 screenings are strains which are perfectly stable and which present the fil phenotype according to the invention. They are resistant to the stress(es) in active metabolism phase and they have still an interesting growth and fermentative power.

The fil strains are intended to be used in industrial applications in which the stress resistance during the active metabolism phase must be high. The properties intended to be obtained for the fil strains are for example a high stability during freezing and/or during drying. The word stability denotes that the fermentative power of the yeast which has been placed under aggressive conditions maintains a high level with respect to the fermentative or gassing power of the same yeast before the aggressive treatment, especially the drying and/or the freezing within a piece of dough.

The stability of the yeast activity of frozen dough after thawing is an important criterion in industrial bakery where the pieces of dough are frozen after beginning of the fermentation, the duration of freezing being from a few days to a few weeks. It is important that the yeast when thawed has not lost the essential part of its fermentative power.

The industrial yeast strain which, on the one hand, has properties which are equivalent to those of industrial strains which are at present marketed in France or in Europe and which, on the other hand, has a stability during freezing in pieces of dough according to test C1 at least higher than 80%, preferably at least higher than 85% and still preferably at least higher than 90%, represents an important progress for the making of frozen doughs. Preferably, this industrial baker's yeast strain is not GMO.

From the point of view of the selection of fil mutants starting from lab strains in order to characterize subsequently the gene(s) which are at the origin of the mutation, or of segregants which are intended to be used for the construction of new industrial strains by classic genetics, experience has shown that it was necessary to adopt at the level of the selection the stability rates which are lower as far as frozen dough is concerned, these strains being in general naturally less resistant to freezing stress.

The baker's yeast, in dried form, presenting at least 92% of dry matter, preferably at least 94% of dry matter, should maintain its fermentative performances, calculated for the same dry matter amount, despite the stress due to dehydration.

It is well known that yeasts present a loss of activity after drying which is all the more lower as they have been harvested during a phase of low growth, that is to say under conditions which are relatively far from those at which they present their maximum fermentative potential. Fil strains allow the shifting of this equilibrium.

That possibility can be measured by comparison of a loss of activity after drying of the *fil* strain with respect to the loss of activity after drying of the starting strain or of a control strain cultured under the same conditions of active growth phase. The loss of activity after drying is defined according to the formula:

$$100 - \frac{A'}{A} \times 100$$

A' represents the fermentative power measured on the dry yeast according to test A', A is the gassing power measured for the yeast before drying according to the corresponding A test.

The process according to the invention has led to the obtaining of two types of mutants of yeast laboratory strains called *fil1* and *fil2*, a different numerical significance (*fil1*, *fil2*,...) being given for each mutation appearing to affect a different gene.

The first family of mutants is the *fil1* family of which the typical strain is the strain PVD1150 = M5 *fil1*. This mutant is derived from the wild type laboratory strain M5 (Schaaff et al., 1989, Curr. Genet., 15, pp.75-81). The *fil1* mutation is carried by the gene *CYR1/CDC35*, which encodes for adenylate cyclase, that is for the enzyme which synthesizes cAMP. The mutation was identified as being a substitution of a glutamate residue by a lysine residue in position 1682 of the protein. This exceptional change is located in the catalytic domain of adenylate cyclase, close to the region which is considered as being implicated in the activation of adenylate cyclase by the Ras proteins. The change of an acidic aminoacid into a basic aminoacid is capable of strongly influencing the activity of the catalytic site, all the more since the change occurs in a very highly conserved zone. This *fil1* mutation introduced in the gene *CYR1/CDC35* of 2 different laboratory strains, caused them to acquire the said *fil* phenotype. Consequently, the *fil1* mutation in the gene *CYR1/CDC35* allows the construction of industrial strains having the *fil* phenotype.

The second family of mutants is the *fil2* family of which the typical strain is the strain KL1 = W303 *fil2*. This mutant is derived from the laboratory strain W303-1A (Thomas et Rothstein (1989) Cell, 56, pp.619-630).

The *fil2* mutation is carried by the gene YDL035c called gene *GPR1* by Xue et al, EMBO J., 1998, 17, 1996-2007. The gene *GPR1* was isolated and identified

marker *URA3*. A laboratory yeast strain auxotrophic for uracil is transformed with this DNA library and, among the transformants which do not need uracil, are selected those which have acquired the *fil* phenotype, by the technique explained concerning the *fil* mutants described above.

- 5 It is remarkable that the different *fil* mutants obtained do not have mutations which affect the same metabolic pathway. This is indicated by the following table:

Properties	Strains			
	<i>fil1</i>	<i>fil2</i>	<i>fil300</i>	AT25 ; AT26 ; AT28 ; AT31
Reduction of the loss of resistance to stress induced by fermentation	yes	yes	yes	yes
Deficiency of cAMP signal induced by glucose	yes	yes	yes	no
Deficiency of accumulation of cAMP induced by acidification	partially yes	no	n.d.	n.d.
Level of trehalose increased	yes	yes	yes	yes
Normal latency phase for culture on glucose	yes	yes	yes	yes
Normal growth rate on glucose	yes	yes	yes	yes
Normal CO ₂ production in the dough	yes	yes	yes	yes
Normal harvest on molasses	yes	yes	yes	yes

n.d.: not determined

10

The increase in cAMP level induced by the glucose is affected in the mutants *fil1*, *fil2* and *fil300*. This is not however the case in the mutants *fil400* (AT25), *fil500* (AT28), and in the mutants AT26 and AT31.

- 15 In the case of the mutants *fil1*, *fil2* and *fil300*, the mutations affect the Ras-cAMP-PKA pathway. This is entirely normal for the *fil1* mutation. This leads to the conclusion that the gene *GPR1*, whose function is not well known, has a function in this pathway.

In the case of the other mutants, the mutation does not seem to concern the Ras-cAMP-PKA pathway.

It is known that the Ras-cAMP-PKA pathway is not the only pathway implicated in the mechanisms of resistance to stress. The new process according to the invention is a tool which is particularly interesting for :

- isolating new mutants resistant to stress
- 5 - characterizing the gene or genes concerned in different metabolic pathways
- constructing with the help of these genes new industrial strains which are particularly efficient.

The resistances to stress are often crossed. The same metabolic pathways are often found in a number of different eukaryotes. In particular, pathways equivalent to the Ras-cAMP-PKA pathway have been described in numerous eukaryotes. Consequently, the present invention is not limited to the obtaining of fresh or dry yeasts for bread-making, for brewing, for winemaking, for alcohol production and distillation, for the production of heterologous proteins, but to the obtaining of all new eukaryotic strains of interest for all industrial uses, such as the production of organic acids, amino-acids, enzymes, etc...

The present invention is also illustrated by the following examples. The list of figures concerning these examples is given hereafter. These figures are designated by the following nomenclature. The first number is the number of the example in which the figure is described, the second number is its order in the said example.

Figure 1-1 : Growth of the strains M5 and M5 *fil1* = PVD1150 on YPD medium at 30°C and under stirring (180 rpm) according to test T3.

Figure 1-2 : cAMP response (Arbitrary Units) of the strains PVD1150 and M5 after addition of glucose (100 mM) to cells cultured on glycerol medium until obtaining of stationary phase.

Figure 1-3 : Monitoring of the degradation of trehalose (Arbitrary Units) in the strains PVD1050, PVD1150 and their respective controls M5 *hxx2Δ* and M5, after an induction by glucose (200 mM) on cells in stationary phase.

Figure 1-4 : Survival rate of the strains HL8.16 *leu2* and HL816 *fil300* after a thermic shock of 30 minutes at 52°C according to test T1.

SUB B1

20

25

30

SUB B2
SUB B3
SUB B4
SUB B5

Figure 1-5 : Survival rate of the strains HL8.16 *leu2* and HL816 *fil300* after a freezing of 12 days at -20°C according to test T2.

Figure 1-6 : Trehalose content (Arbitrary Units) in the strains HL8.16 *leu2* and HL816 *fil300*. The strains are grown until obtaining of stationary phase, and then glucose (100 mM) is added at t=0.

Figure 1-7 : cAMP response (Arbitrary Units) after induction by glucose (100 mM) in the strains HL8.16 *leu2* and HL816 *fil300* cultured until obtaining of stationary phase. The initial adding of 3 mM of glucose allows to avoid the cAMP response related to intracellular acidification.

Figure 2-1 : Monitoring of the degradation of trehalose (Arbitrary Units) after an induction by glucose at t = 0 (on cells of strains S47, AT25, AT26, AT28, AT31 in stationary phase).

Figure 2-2 : Monitoring of the cAMP response (Arbitrary Units) after an induction (at t = 0) by glucose on cells in exponential growth phase on maltose. a) control S47 and mutant AT25 ; b) control S47 and mutants AT26, AT28 and AT31.

Figure 3-1 : Gap-filling strategy (filling of missing DNA) used for isolating the gene carrying the *fil1* mutation in the strain PVD1150.

Figure 3-2 : Physical map of the vector pUC18-CYR1mut -URA3 [Sn].

Figure 6-1 : Stability to freezing measured by the ratio between the gassing power after 1 day of conservation at -20°C and n days of storage at -20°C.

EXAMPLE 1 - Use of heat stress for the isolation of fil mutants

A. Obtaining of strains having the *fil1* phenotype

a) Obtaining of the strains PVD1050 and PVD1150

The starting strain is the haploid strain M5 *hvk2Δ* (strain comprising a deletion of the gene of hexokinase II) which is derived from the haploid strain M5, which is derived from the diploid strain M5 (Schaaff et coll. (1989), Curr. Genet. 15: 75-81). A mutagenesis using E.M.S. (Ethyl Methyl Sulfonate) has been carried out according to the technique disclosed in "Methods in Yeast Genetics, Cold Spring Harbor Laboratory Press" (Sherman et coll., 1986) in such a way as to obtain a survival rate of about 10%. After this treatment, the cells are washed and resuspended

in the YPD medium. They are then cultivated until stationary phase is reached. A 0.25 ml sample of this culture has been taken and used for the inoculation of 25 ml of YPD medium. The inoculated medium has been incubated at 30°C for 90 minutes, and then heated to 52°C for 30 minutes. It has been then incubated at 30°C for 24 hours, and then 0.5 ml of the culture were transferred into 25 ml of YPD medium. This medium has been incubated at 30°C for 90 minutes. Fractions of 100 µl of this medium were then taken and placed at 56°C for 30 minutes. Then, the cell suspension was spread on YPD-A medium. The stress resistance was then evaluated by tests on surviving cells forming colony on YPD-A medium.

In order to test the said resistance, the yeast cells of the strains as obtained were tested according to the test T1, using an incubation at 52°C for 30 minutes. Under the conditions of this test and after verification of the conservation of the growth properties and of the fermentation properties, as well as of the stability of the mutation, one strain has been selected and called Mut1 or PVD1050. It has been shown that this strain comprises a stable mutation, which was called *fill* and which is monogenic and recessive. The haploid strain PVD1050 has a survival rate after thermic treatment which is surprisingly high, and this high survival rate is observed for stationary cells and also for cells growing in exponential phase, as shown by the results collected in table 1-A. It is surprising that, in the case of a *Saccharomyces cerevisiae* strain, 100% viability is conserved in exponential phase after a thermic treatment of 30 minutes at 52°C. It is also surprising to notice that this strain PVD1050 is not significantly penalized from the standpoint of growth and fermentation with respect to the strain M5 *hvk2Δ*, being recalled that the deletion of the gene *HVK2* penalizes among others the growth.

In order to eliminate the *HVK2* deletion from this strain, the haploid strain PVD1050 (=M5 *hvk2Δ fill*) has been crossed with the haploid strain M5 and subsequently tetrads were dissected. In the thus obtained haploids, a strain presenting the *fill* mutation but having also a wild type *HVK2* gene was selected. The thus isolated haploid strain was called PVD1150 or M5 *fill*. That strain has the same thermoresistance phenotype as the strain PVD1050, i.e. a survival rate which is extremely high in the test T1 (table 1-A).

EVALUATION OF THE SURVIVAL RATE OF *fill* STRAINS AND OF THEIR CONTROLS AFTER A THERMIC SHOCK OF 30 MINUTES AT 52°C

Test T1 with a survival rate after 30' at 52°C (%)				
	In stationary Phase	In active metabolism = glucose incubation of:		
		30'	60'	90'
M5 <i>hxx2Δ</i>	70 %	40 %	30 %	15 %
PVD1050	100 %	98 %	98 %	98 %
M5	50 %	20 %	10 %	5 %
PVD1150	99 %	96 %	-	-

5

10

15

20

Table 1-B

	Test T5 Harvest (expressed in dry matter of yeast) after 40 h [with respect to 8 g of molasses]	Test A20 Fermentative ability (CO ₂ ml released in 2 hours)
M5	1.70 g	66-75 ml
M5 <i>fill</i>	1.74 g	60-65 ml

From this table, it appears that there is no penalization during growth and a low penalization or decrease of the fermentation as far as the mutant is concerned. It results that the mutant is a mutant which presents very likely the new *fil* phenotype, object of the invention.

In the case of the two strains presenting the *fill* mutation (PVD1050 and PVD1150) the levels of cAMP and of trehalose were examined with respect to their respective controls (M5 *hxx2Δ* and M5). The cAMP level has been measured on the strains PVD1150 and M5, according the method disclosed by Thévelein et coll., 1987, J. Gen. Microbiol., 133, pp.2197-2205. The cAMP level has thus been determined after induction of its synthesis by addition of glucose (100 mM) on cells which have reached the stationary phase after growth on glycerol. An attenuated cAMP signal has been shown in the mutant PVD1150 (figure 1-2). Trehalose has also been determined on the strains PVD1150, PVD1050, M5 and M5 *hxx2Δ*, according to the conditions described by Neves et coll., 1991, FEBS Lett., 283, pp.19-22. In the case of strains presenting the *fill* mutation, the mobilization of the trehalose (after induction by glucose on cells in stationary phase) is far less rapid than in the control strains (figure 1-3).

From the results of the determination of trehalose and of cAMP, it seems obvious that the *fill* mutation concerns the Ras-cAMP pathway. However, and in contradiction with respect to the mutants of the Ras-cAMP pathway previously isolated and which show growth delays and fermentation delays, the *fill* mutant is only slightly affected during growth and fermentation.

In order to determine the actual influence of the increase trehalose content on the stress resistance of mutants *fill*, the gene *TPS1* (which codes for the synthesis of trehalose phosphate starting from glucose) has been deleted in the strains *fill* and in the

6674
 6675
 6676
 6677
 6678
 6679
 6680
 6681
 6682
 6683
 6684
 6685
 6686
 6687
 6688
 6689
 6690
 6691
 6692
 6693
 6694
 6695
 6696
 6697
 6698
 6699
 6700
 6701
 6702
 6703
 6704
 6705
 6706
 6707
 6708
 6709
 6710
 6711
 6712
 6713
 6714
 6715
 6716
 6717
 6718
 6719
 6720
 6721
 6722
 6723
 6724
 6725
 6726
 6727
 6728
 6729
 6730
 6731
 6732
 6733
 6734
 6735
 6736
 6737
 6738
 6739
 6740
 6741
 6742
 6743
 6744
 6745
 6746
 6747
 6748
 6749
 6750
 6751
 6752
 6753
 6754
 6755
 6756
 6757
 6758
 6759
 6760
 6761
 6762
 6763
 6764
 6765
 6766
 6767
 6768
 6769
 6770
 6771
 6772
 6773
 6774
 6775
 6776
 6777
 6778
 6779
 6780
 6781
 6782
 6783
 6784
 6785
 6786
 6787
 6788
 6789
 6790
 6791
 6792
 6793
 6794
 6795
 6796
 6797
 6798
 6799
 6800
 6801
 6802
 6803
 6804
 6805
 6806
 6807
 6808
 6809
 6810
 6811
 6812
 6813
 6814
 6815
 6816
 6817
 6818
 6819
 6820
 6821
 6822
 6823
 6824
 6825
 6826
 6827
 6828
 6829
 6830
 6831
 6832
 6833
 6834
 6835
 6836
 6837
 6838
 6839
 6840
 6841
 6842
 6843
 6844
 6845
 6846
 6847
 6848
 6849
 6850
 6851
 6852
 6853
 6854
 6855
 6856
 6857
 6858
 6859
 6860
 6861
 6862
 6863
 6864
 6865
 6866
 6867
 6868
 6869
 6870
 6871
 6872
 6873
 6874
 6875
 6876
 6877
 6878
 6879
 6880
 6881
 6882
 6883
 6884
 6885
 6886
 6887
 6888
 6889
 6890
 6891
 6892
 6893
 6894
 6895
 6896
 6897
 6898
 6899
 6900
 6901
 6902
 6903
 6904
 6905
 6906
 6907
 6908
 6909
 6910
 6911
 6912
 6913
 6914
 6915
 6916
 6917
 6918
 6919
 6920
 6921
 6922
 6923
 6924
 6925
 6926
 6927
 6928
 6929
 6930
 6931
 6932
 6933
 6934
 6935
 6936
 6937
 6938
 6939
 6940
 6941
 6942
 6943
 6944
 6945
 6946
 6947
 6948
 6949
 6950
 6951
 6952
 6953
 6954
 6955
 6956
 6957
 6958
 6959
 6960
 6961
 6962
 6963
 6964
 6965
 6966
 6967
 6968
 6969
 6970
 6971
 6972
 6973
 6974
 6975
 6976
 6977
 6978
 6979
 6980
 6981
 6982
 6983
 6984
 6985
 6986
 6987
 6988
 6989
 6990
 6991
 6992
 6993
 6994
 6995
 6996
 6997
 6998
 6999
 7000
 7001
 7002
 7003
 7004
 7005
 7006
 7007
 7008
 7009
 7010
 7011
 7012
 7013
 7014
 7015
 7016
 7017
 7018
 7019
 7020
 7021
 7022
 7023
 7024
 7025
 7026
 7027
 7028
 7029
 7030
 7031
 7032
 7033
 7034
 7035
 7036
 7037
 7038
 7039
 7040
 7041
 7042
 7043
 7044
 7045
 7046
 7047
 7048
 7049
 7050
 7051
 7052
 7053
 7054
 7055
 7056
 7057
 7058
 7059
 7060
 7061
 7062
 7063
 7064
 7065
 7066
 7067
 7068
 7069
 7070
 7071
 7072
 7073
 7074
 7075
 7076
 7077
 7078
 7079
 7080
 7081
 7082
 7083
 7084
 7085
 7086
 7087
 7088
 7089
 7090
 7091
 7092
 7093
 7094
 7095
 7096
 7097
 7098
 7099
 7100
 7101
 7102
 7103
 7104
 7105
 7106
 7107
 7108
 7109
 7110
 7111
 7112
 7113
 7114
 7115
 7116
 7117
 7118
 7119
 7120
 7121
 7122
 7123
 7124
 7125
 7126
 7127
 7128
 7129
 7130
 7131
 7132
 7133
 7134
 7135
 7136
 7137
 7138
 7139
 7140
 7141
 7142
 7143
 7144
 7145
 7146
 7147
 7148
 7149
 7150
 7151
 7152
 7153
 7154
 7155
 7156
 7157
 7158
 7159
 7160
 7161
 7162
 7163
 7164
 7165
 7166
 7167
 7168
 7169
 7170
 7171
 7172
 7173
 7174
 7175
 7176
 7177
 7178
 7179
 7180
 7181
 7182
 7183
 7184
 7185
 7186
 7187
 7188
 7189
 7190
 7191
 7192
 7193
 7194
 7195
 7196
 7197
 7198
 7199
 7200
 7201
 7202
 7203
 7204
 7205
 7206
 7207
 7208
 7209
 7210
 7211
 7212
 7213
 7214
 7215
 7216
 7217
 7218
 7219
 7220
 7221
 7222
 7223
 7224
 7225
 7226
 7227
 7228
 7229
 7230
 7231
 7232
 7233
 7234
 7235
 7236
 7237
 7238
 7239
 7240
 7241
 7242
 7243
 7244
 7245
 7246
 7247
 7248
 7249
 7250
 7251
 7252
 7253
 7254
 7255
 7256
 7257
 7258
 7259
 7260
 7261
 7262
 7263
 7264
 7265
 7266
 7267
 7268
 7269
 7270
 7271
 7272
 7273
 7274
 7275
 7276
 7277
 7278
 7279
 7280
 7281
 7282
 7283
 7284
 7285
 7286
 7287
 7288
 7289
 7290
 7291
 7292
 7293
 7294
 7295
 7296
 7297
 7298
 7299
 7300
 7301
 7302
 7303
 7304
 7305
 7306
 7307
 7308
 7309
 7310
 7311
 7312
 7313
 7314
 7315
 7316
 7317
 7318
 7319
 7320
 7321
 7322
 7323
 7324
 7325
 7326
 7327
 7328
 7329
 7330
 7331
 7332
 7333
 7334
 7335
 7336
 7337
 7338
 7339
 7340
 7341
 7342
 7343
 7344
 7345
 7346
 7347
 7348
 7349
 7350
 7351
 7352
 7353
 7354
 7355
 7356
 7357
 7358
 7359
 7360
 7361
 7362
 7363
 7364
 7365
 7366
 7367
 7368
 7369
 7370
 7371
 7372
 7373
 7374
 7375
 7376
 7377
 7378
 7379
 7380
 7381
 7382
 7383
 7384
 7385
 7386
 7387
 7388
 7389
 7390
 7391
 7392
 7393
 7394
 7395
 7396
 7397
 7398
 7399
 7400
 7401
 7402
 7403
 7404
 7405
 7406
 7407
 7408
 7409
 7410
 7411
 7412
 7413
 7414
 7415
 7416
 7417
 7418
 7419
 7420
 7421
 7422
 7423
 7424
 7425
 7426
 7427
 7428
 7429
 7430
 7431
 7432
 7433
 7434
 7435
 7436
 7437
 7438
 7439
 7440
 7441
 7442
 7443
 7444
 7445
 7446
 7447
 7448
 7449
 7450
 7451
 7452
 7453
 7454
 7455
 7456
 7457
 7458
 7459
 7460
 7461
 7462
 7463
 7464
 7465
 7466
 7467
 7468
 7469
 7470
 7471
 7472
 7473
 7474
 7475
 7476
 7477
 7478
 7479
 7480
 7481
 7482
 7483
 7484
 7485
 7486
 7487
 7488
 7489
 7490
 7491
 7492
 7493
 7494
 7495
 7496
 7497
 7498
 7499
 7500
 7501
 7502
 7503
 7504
 7505
 7506
 7507
 7508
 7509
 7510
 7511
 7512
 7513
 7514
 7515
 7516
 7517
 7518
 7519
 7520
 7521
 7522
 7523
 7524
 7525
 7526
 7527
 7528
 7529
 7530
 7531
 7532
 7533
 7534
 7535
 7536
 7537
 7538
 7539
 7540
 7541
 7542
 7543
 7544
 7545
 7546
 7547
 7548
 7549
 7550
 7551
 7552
 7553
 7554
 7555
 7556
 7557
 7558
 7559
 7560
 7561
 7562
 7563
 7564
 7565
 7566
 7567
 7568
 7569
 7570
 7571
 7572
 7573
 7574
 7575
 7576
 7577
 7578
 7579
 7580
 7581
 7582
 7583
 7584
 7585
 7586
 7587
 7588
 7589
 7590
 7591
 7592
 7593
 7594
 7595
 7596
 7597
 7598
 7599
 7600
 7601
 7602
 7603
 7604
 7605
 7606
 7607
 7608
 7609
 7610
 7611
 7612
 7613
 7614
 7615
 7616
 7617
 7618
 7619
 7620
 7621
 7622
 7623
 7624
 7625
 7626
 7627
 7628
 7629
 7630
 7631
 7632
 7633
 7634
 7635
 7636
 7637
 7638
 7639
 7640
 7641
 7642
 7643
 7644
 7645
 7646
 7647
 7648
 7649
 7650
 7651
 7652
 7653
 7654
 7655
 7656
 7657
 7658
 7659
 7660
 7661
 7662
 7663
 7664
 7665
 7666
 7667
 7668
 7669
 7670
 7671
 7672
 7673
 7674
 7675
 7676
 7677
 7678
 7679
 7680
 7681
 7682
 7683
 7684
 7685
 7686
 7687
 7688
 7689
 7690
 7691
 7692
 7693
 7694
 7695
 7696
 7697
 7698
 7699
 7700
 7701
 7702
 7703
 7704
 7705
 7706
 7707
 7708
 7709
 7710
 7711
 7712
 7713
 7714
 7715
 7716
 7717
 7718
 7719
 7720
 7721
 7722
 7723
 7724
 7725
 7726
 7727
 7728
 7729
 7730
 7731
 7732
 7733
 7734
 7735
 7736
 7737
 7738
 7739
 7740
 7741
 7742
 7743
 7744
 7745
 7746
 7747
 7748
 7749
 7750
 7751
 7752
 7753
 7754
 7755
 7756
 7757
 7758
 7759
 7760
 7761
 7762
 7763
 7764
 7765
 7766
 7767
 7768
 7769
 7770
 7771
 7772
 7773
 7774
 7775
 7776
 7777
 7778
 7779
 7780
 7781
 7782
 7783
 7784
 7785
 7786
 7787
 7788
 7789
 7790
 7791
 7792
 7793
 7794
 7795
 7796
 7797
 7798
 7799
 7800
 7801
 7802
 7803
 7804
 7805
 7806
 7807
 7808
 7809
 7810
 7811
 7812
 7813
 7814
 7815
 7816
 7817
 7818
 7819
 7820
 7821
 7822
 7823
 7824
 7825
 7826
 7827
 7828
 7829
 7830
 7831
 7832
 7833
 7834
 7835
 7836
 7837
 7838
 7839
 7840
 7841
 7842
 7843
 7844
 7845
 7846
 7847
 7848
 7849
 7850
 7851
 7852
 7853
 7854
 7855
 7856
 7857
 7858
 7859
 7860
 7861
 7862
 7863
 7864
 7865
 7866
 7867
 7868
 7869
 7870
 7871
 7872
 7873
 7874
 7875
 7876
 7877
 7878
 7879
 7880
 7881
 7882
 7883
 7884
 7885
 7886
 7887
 7888
 7889
 7890
 7891
 7892
 7893
 7894
 7895
 7896
 7897
 7898
 7899
 7900
 7901
 7902
 7903
 7904
 7905
 7906
 7907
 7908
 7909
 7910
 7911
 7912
 7913
 7914
 7915
 7916
 7917
 7918
 7919
 7920
 7921
 7922
 7923
 7924
 7925
 7926
 7927
 7928
 7929
 7930
 7931
 7932
 7933
 7934
 7935
 7936
 7937
 7938
 7939
 7940
 79

starting control strains. Despite the fact that these strains *fil1 tps1Δ* no longer synthesize trehalose, they retain a much better resistance against thermic stress than the control strains *tps1Δ* : 30% survival after 30 minutes at 52°C, the survival being 0.1% as far as the controls are concerned (conditions of test T1). This confirms the results published by Van Dijck et coll. (1995, Appl. Environ. Microbiol., 61, pp.109-115) which demonstrated the fact that trehalose is not the sole responsible for the improvement of stress resistance.

B. Isolation of strains *fil300*

a/ Isolation of the strain FD51 = HL816 *fil300*

A mutagenesis has been carried out on strain HL8.16 *leu2* (aneuploid strain, segregant of an industrial strain made auxotrophic for leucine ; LESAFFRE's collection) : the cells have been cultured at 30°C in a YPD medium until the stationary phase was obtained. Then the cells have been incubated at 30°C for 1 hour, in the presence of E.M.S. (Ethyl Methyl Sulfonate) according to the technique disclosed in " Methods in Yeast Genetics, Cold Spring Harbor Laboratory Press " (Sherman et al., 1986) in such a way that a survival rate of about 10% is obtained. After that treatment, the cells have been washed, resuspended in YPD medium and cultured at 30°C until the stationary phase was obtained. Glucose has then been added in the culture in stationary phase in order to obtain a final concentration in glucose equal to 100 mM. The inoculated medium has been incubated at 30°C for 30 minutes, then at 56°C for 30 minutes, and finally at 65°C for 30 minutes. The surviving cells have been isolated by spreading on YPD-A medium. Then they have been subjected individually to a test of resistance to heat, according to the conditions of test T1 with an incubation of 30 minutes at 52°C. At the end of this first selection, the six strains which presented the highest survival rates have been retained and subjected to a second mutagenic treatment, with ultraviolet rays (U.V.), according to the technique disclosed by Spencer J.F.T. and Spencer D.M., Chapter " Yeast genetics " extract of " Yeast a practical approach ", 1988, Campbell and Duffus Eds. For this treatment, the cells of each of the six strains have been cultivated in a YPD medium until obtaining of the stationary phase. They were then washed with water, diluted and spread on YPD-A medium. U.V. light at a dose of 30 mJ (wavelength equal to 260 nm) has been applied on the

open dishes. The 248 colonies which survived the mutagenic treatment were transferred to a YPD-A medium and then incubated during 2 hours at 60°C. At the end of this thermic treatment, only 2 colonies survived. A test of resistance to heat in liquid medium has then been carried out on these two colonies, according to the conditions of test T1 with an incubation of 30 minutes at 52°C after an incubation in the presence of glucose for 1 hour at 30°C. The best of the 2 strains was subjected to a last mutagenic treatment with U.V. light (ultraviolet rays), according to the conditions previously disclosed however with a dose of 10 mJ of U.V. light (260 nm). The colonies which survived the mutagenic treatment were transferred to a YPD-A medium and incubated during 6 hours at 60°C. At the end of this thermic treatment, 171 colonies (of close to 1500) survived. A heat resistance test carried out on these 171 strains according to the conditions of test T1 permitted the selection of strain FD51, also called HL816 *fil300*. It has of course been checked that the thus obtained mutation is stable and that it corresponds to all of the conditions of *fil* phenotype.

b/ Characterization of the strain HL816 *fil300*

A significant improvement of the heat resistance of the strain has been confirmed by test T1, either in stationary phase or in active fermentation phase (figure 1-4). Furthermore, mutation *fil300* induced a resistance against other stresses : strain HL816 *fil300* presents, under conditions of test T2, a survival rate of more than 50% after freezing during 12 days at -20°C of pre-fermented cells during 90 minutes at 30°C while the survival rate of the control is in the same conditions lower than 11% (figure 1-5). The strain HL816 *fil300* presents consequently a much better resistance against freezing than the control strain, whether the cells are in stationary phase or in active fermentation phase.

In order to better characterize the phenotype linked to the *fil300* mutation, the levels of trehalose and cAMP have been determined on cells in stationary phase which were subjected to an induction by glucose. The control strain is the starting strain HL8.16 *leu2*. During stationary phase (i.e. 0 minute of fermentation in figure 1-6), the trehalose level is 3 to 4 times higher in mutant *fil300* than in control HL8.16 *leu2*. During exponential phase, the degradation rate of trehalose in the mutant HL8.16 *fil300* strain is reduced, with respect to that of the control (figure 1-6). In mutant

09330265-06110

SUB
B16

SUB
B17

SUB
B17
COO-t.

HL816 *fil300* a cAMP response reduced by close to 50% after an induction of the synthesis by glucose on cells in stationary phase has also been observed (figure 1.7).

The performances of growth and of fermentative ability of strains HL8.16 *leu2* (control) and HL816 *fil300* were determined according to the tests T5 and A20. The results are presented in table 1-C. The strain *fil300* retains in these tests 80% of the characteristic features measured for the starting strain.

Table 1-C

	T5	A20
	Harvest (in dry matter of yeast) 20 h [for 8 g of molasses]	Fermentative ability (ml CO ₂ released in 2 hours)
HL8.16 <i>leu2</i>	1.5 g	85
HL816 <i>fil300</i>	1.2 g	68

b/ Isolation of the strain HL822 *fil300*

The decreases of biomass growth yield and of fermentative ability of mutant HL816 *fil300* which are at the maximum of what is tolerated within the frame of *fil* phenotype are detrimental to its utilization. This is the reason why crossings have been carried out between the said mutant and some segregants in order to isolate in the descendance of the diploids obtained, thermoresistant segregants having improved performances. The strain HL822 *fil300*, also called FDH16-22, has thus been isolated. This strain, which is a segregant issued from a crossing between the strain HL816 *fil300* and the segregant HL816 (LESAFFRE's collection), has a growth yield on molasse and a gassing power identical to those of the control strain HL8.16 *leu2* (table 1-D). It presents furthermore a level of thermoresistance identical to that of the strain HL816 *fil300*.

Table 1-D

GROWTH YIELD ON MOLASSES AND FERMENTATIVE ABILITY (DETERMINED
IN TEST A20) OF THE STRAINS HL8.16 LEU2 AND HL822 *FIL300*

	T5	A20
	Harvest (in dry matter of yeast) after 20 h [for 8 g of molasses]	Fermentative ability (ml CO ₂ released in 2 hours)
HL8.16 leu2	1.5 g	85
HL822 <i>fil300</i>	1.5 g	85

5

With respect to each of the two *fil300* strains (HL816 *fil300* and HL822 *fil300*), the stability under the form of frozen doughs has been compared with that of the control strain HL8.12 *leu2* by monitoring the gassing power as a function of freezing duration under the conditions of test C2.

10

A significant improvement of the stability during freezing has been observed for the *fil300* strains with respect to the control HL8.16 *leu2* (table 1-E). Thus, the *fil300* strains retain at least 60% of their fermentative ability (= gassing power) after 1 month of conservation at the temperature of -20°C while the control only conserves 40% of its gassing power under the same conditions (table 1-E).

15

Table 1-E

STABILITY DURING FREEZING, DETERMINED BY THE RATIO BETWEEN
THE FERMENTATIVE ABILITY AT DAY N OF MEASUREMENT AND
THE FERMENTATIVE ABILITY AFTER THE FIRST DAY OF FREEZING (TEST C2)

20

	Test C2		
	Stability during freezing (%)		
	1 day	28 days	42 days
HL8.16 leu2	100 %	41 %	39 %
HL816 <i>fil300</i>	100 %	60 %	53 %
HL822 <i>fil300</i>	100 %	62 %	57 %

EXAMPLE 2 - Use of the cycles of freezing/thawing for isolation of fil mutants starting from an industrial strain

The starting or original strain is the strain S47 deposited at the C.N.C.M.,
 5 25 rue de Docteur Roux, 75724 Paris, France, under the number I-2037. This strain is
 an aneuploid baker's yeast strain used industrially, it is probably the best strain at
 present used in France for making raw frozen doughs. It has been grown in a liquid
 YPD medium at 30°C until obtaining of the stationary phase. A mutagenic treatment
 with U.V. was carried out according to the technique described by Spencer J.F.T. and
 10 Spencer D.M., Chapter "Yeast genetics" extract of "Yeast a practical approach"
 (1988), Campbell and Duffus Eds., in order to obtain a survival rate of about 10%.
 The cells have thus been treated with a dose of U.V. (260 nm) of 5 mJ. They have then
 been grown at 30°C on a medium consisting of agar YE-molasses (agar 20 g/l, extract
 of yeast 5 g/l, molasses 5 g/l, $(\text{NH}_4)_2\text{HPO}_4$ 0.5 g/l, pH 5.0-5.5), and were then
 15 harvested after 2 to 3 days by washing of the dishes with water. Mini-pieces of dough
 of 0.5 g were prepared with the harvested yeasts, with the following composition :
 flour 56.5 %, NaCl 1.0 %, water 42.5 %, yeasts 4.10^8 cells/gram of dough. These mini-
 pieces of dough were incubated at 30°C for 30 minutes. Then they were subjected to
 200 successive cycles of freezing at -30°C and thawing at room (ambient) temperature,
 20 in order to obtain only a few thousand surviving cells (survival rate lower than 0.01%),
 which were isolated from the pieces of doughs.

The freeze resistance has then been tested individually on the surviving
 cells, by determination of survival rate of the cells after one freezing and by
 determination of the loss of fermentative ability (= gassing power) after freezing. In
 25 parallel, the growth yield on molasses and the fermentative ability (= gassing power) of
 the best strains have been measured.

- * The survival rate after freezing is measured according to the conditions of test T2
 with a fermenting phase of 30 minutes at 30°C, followed by a freezing at -30°C for 24
 hours, and then by a thawing at room temperature.
- 30 * The growth of the strains has been measured by the yield obtained after a culture for
 20 hours on agar molasses medium according to test T5 ; the harvested yeasts are used

to determine the fermentative ability at 30°C and over 2 hours, according to the conditions of test A1.

From the strains which survived the successive treatments of freezing/thawing in mini-pieces of dough, 7 strains named AT25, AT26, AT27, AT28, AT29, AT30 and AT31 were pre-selected. The results of the survival rates after freezing (test T2) are collected in table 2-A, the molasses growth yields (T5) and the gassing power (A1) are collected in table 2-B. Even if all of these mutants present a survival rate (table 2-A) much better than that of the control, it is necessary to take into account the growth yield and gassing power of each mutant for the selection of strains which correspond to criteria of fil phenotype (table 2-B). This is the reason why only the strains AT25, AT26, AT28 and AT31 are selected at the end of these tests. The measurements of the maximal growth rate were carried out according to test T4 on these different mutants and the control strain S47, the results being indicated in table 2-C.

Table 2-A

	Test T2	
	Survival rate (%) after :	
	Freezing without previous fermentation -25°C during 24 hours	Freezing (-25°C during 24 hours) with previous fermentation of 30 minutes at 30°C
S47	36	17
AT25	51	47
AT26	59	42
AT27	45	40
AT28	47	39
AT29	72	32
AT30	47	28
AT31	38	27

Table 2-B

GROWTH YIELD (EXPRESSED BY THE RATIO BETWEEN THE AMOUNT OF YEAST PRODUCED IN 20 HOURS EXPRESSED AT 30% OF DRY MATTER WITH RESPECT TO THE AMOUNT OF MOLASSES USED CONSIDERED AS BEING 50% OF SUGAR) AND FERMENTATIVE ABILITY (EXPRESSED BY THE AMOUNT OF CO₂ RELEASED IN 2 HOURS AT 30°C) OF THE MUTANTS OF THE STRAIN S47 AND OF THE STARTING STRAIN S47

	Test T5 Growth yield (%)	Test A1 Fermentative ability (ml CO ₂)
S47	67 %	138
AT25	64 %	116
AT26	62 %	116
AT27	47 %	113
AT28	62 %	120
AT29	51 %	64
AT30	62 %	84
AT31	62 %	133

Table 2-C

10 MAXIMAL GROWTH RATE OF STRAIN S47 AND OF THE ISOLATED MUTANTS DERIVED FROM STRAIN S47, ON DIFFERENT GROWTH MEDIA ACCORDING TO TEST T4

STRAIN	MAXIMAL GROWTH RATE or μ_{\max} (h ⁻¹)		
	glucose 100 mM	glucose 10 mM	molasse 0.5%
S47	0.69	0.60	0.57
AT25	0.60	0.61	0.54
AT26	0.67	0.56	0.56
AT27	0.37	0.23	0.26
AT28	0.61	0.62	0.60
AT29	0.59	0.48	0.52
AT30	0.62	0.55	0.56
AT31	0.63	0.60	0.57

In order to check whether the mutations carried by the strains AT25, AT26, AT28 and AT31 also affect the Ras-cAMP pathway as in the case of the *fil* strains described in example 1, the levels of cAMP and of trehalose in these different strains have been measured. The measurements of trehalose have been carried out on cells

 5 cultured until the stationary phase and then subjected to an induction of its mobilization or degradation by addition of glucose (100 mM). The initial level of trehalose, which corresponds to the level of trehalose in stationary phase, is higher for all the mutants than for the control ($t=0$, figure 2-1). In the presence of glucose, the degradation of trehalose is rather quick for each of the said strains (mutants and

 10 control) as after 20 minutes and more, a low and almost identical level of trehalose is obtained for all of these strains. However, the stress imposed within the frame of test T2 is realized after an incubation of 30 minutes in the presence of glucose. This confirms, here again, that trehalose does not allow an explanation of the better performances of these *fil* mutants. As far as cAMP is concerned, the measurements

 15 were carried out on cells which were in exponential growth phase on maltose which have been subjected to an induction by glucose (100 mM). In these experiments, there is no significant reduction of the cAMP signal in the mutants with respect to the control strains S47 (figures 2-2, a & b). Consequently, it seems that the cryoresistant mutants AT25, AT26, AT28 and AT31, which had been isolated following successive

 20 cycles of freezing/thawing and which have the *fil* phenotype, carry mutations which affect other targets than the Ras-cAMP pathway.

25 **EXAMPLE 3 : Identification of the *fil* mutation and reconstruction of strains carrying the *fil* mutation**

Crossings between the strain PVD1150 and other strains have shown that the *fil* mutation is located close to a centromere. The strain PVD1150 has consequently been recrossed with different haploid strains carrying genetic markers

 30 located close to the centromeres of each chromosome and analysis has been carried out on the tetrads originating from the sporulation of the thus obtained diploids.

SUB
B18

657F30 23202260

THE
NEW
YORK
PUBLIC
LIBRARY
ASTOR LENOX TILDEN FOUNDATION

- 15

20

1/ Transformation of the strain PVD1150 by centromeric plasmids carrying each one of the genes located at less than 25 kb of the centromere of chromosome X

25

30

2/ Search for the transformants having lost the phenotype of thermoresistance

The different transformants carrying each, on a monocopy vector, one of the genes located at least at 25 kb from the centromere of chromosome X, have been tested for their loss of thermoresistance. Each strain PVD1150 carrying one of the genes located around the centromere of chromosome X is streaked on YPD-A medium preheated at 57°C, which is incubated for 90 minutes at 57°C. In parallel, the strain PVD1150 which contains a monocopy plasmid without insert, here the vector YCplac33, is used as non complemented control. After this thermic treatment, the strains are incubated at 30°C until development of the growth of the control (appearance of a cellular layer). The strains carrying a gene which complements the *fil1* mutation are those which have not yet developed.

The results of these tests allowed to demonstrate that the *fil1* mutation is located in the gene *CYR1* (also called *CDC35*), as only that gene was capable of complementing the *fil* phenotype in strain PVD1150. The gene *CYR1/CDC35* codes for adenylate cyclase, enzyme of the Ras-cAMP pathway which permits the synthesis of cAMP starting from ATP. This result is coherent with the fact that the PVD1150 strain presents a reduced level of cAMP (example 1).

In order to check that the loss of thermoresistance is actually linked to the presence of the wild type gene *CYR1/CDC35* introduced by the plasmid, the strain PVD1150 containing the vector YCplac33-CYR1 has been cultured on YPD medium, a medium which is non selective and which permits rapid loss of the plasmid. Colonies have been isolated from that culture on a YPD-A medium, and then replicated on selective medium SD-URA (medium deprived of uracil) in order to search for clones which do not grow and which consequently have lost the plasmid. A thermoresistance test identical to the previous one confirms that all these clones having lost the plasmid had again become thermoresistant.

3/ Isolation of the gene carrying the mutation from the mutant PVD1150

The deletion of the gene is known as having as consequence the lethality of the strains. It was necessary to isolate the mutated gene.

The mutated gene of the strain PVD1150 has been cloned by the technique of “ gap-filling ” [Iwasaki, T. and coll. (1991) *Gene*, 109, pp.81-87] or “ allele rescue ” [Orr-Weaver, T.L. et coll. (1983) “ *Methods Enzymol.* ”, 101, pp.228-245].

This method has been used with the strain PVD1150 by transformation with the plasmid YCplac33-CYR1 digested by the enzyme *Sna*BI (figure 3-1). The transformants which grow on minimum medium without uracil (medium SD-URA) have then been selected. These transformants can only grow when there has been a recircularization by the vector, especially coming from an event of double recombination having integrated the missing part of the gene due to the mutated gene *CYR1* in the PVD1150 strain (figure 3-1). In parallel, the strain M5 has been transformed with the vector YCplac33-CYR1 digested by *Sna*BI, then the transformants have been selected on minimum medium in order to obtain a vector carrying the non mutated gene *CYR1* originated from the control strain M5.

The strain PVD1150 has then been retransformed :

- with the vector YCplac33-CYR1^{mut} which carries the mutated gene originating from vector YCplac33-CYR1 digested by *Sna*BI and filled by gap-filling in the strain PVD1150,
- with the vector YCplac33-CYR1/S which carries the wild type gene originating from vector YCplac33-CYR1 digested by *Sna*BI and filled by gap-filling in the strain M5.

The level of thermoresistance of these two strains has then been compared with the help of the following test : streaks of cells YPD-A medium preheated at 57°C and incubation at 57°C for 90 minutes. As expected, the mutated gene does not complement the mutation ; on the other hand, the wild type *CYR1* gene makes the strain PVD1150 thermosensitive.

4/ Sequencing of the mutated gene (originating from PVD1150) and of its wild type allele (originating from M5) in order to identify and locate the mutation

The mutated gene *CYR1*^{mut} originating from PVD1150 and the wild type *CYR1* gene originating from M5 were sequenced in parallel by the technique of direct sequencing on PCR products obtained by amplification of fragments covering the whole gene, due to which it became possible to locate the *fil1* mutation. It consists of the change of one base G into one base A in position n° 5044 of the coding part of gene *CYR1* (gene referenced under the name YJL005w in the case of *Saccharomyces*

cerevisiae).

This also corresponds to a change in position 429888 of chromosome X, according to the classification of MIPS (Munich Information Center for Protein Sequence). This change is represented in the following table.


5

<u>Control strain</u>	<u><i>fil</i> strain</u>
glu	lys
GGGAGCT	GGAAGCT
CCCTC GA	CCTTC GA

10

The mutation has been verified both by sequencing of a mixture of three independent PCR products and by sequencing of the complementary DNA strand. This change leads to the change from a glutamic acid into lysine in position 1682 of the protein. It occurs in the region coding for the catalytic site of the enzyme, and close to the region which is supposed to be involved in the activation of adenylate cyclase by the Ras proteins :

15



20

5/ Reintroduction (by homologous recombination) of the mutated gene in the thermosensitive lab strains and characterization of the associated phenotype

a/ Construction of the plasmid pUC18-CYR1^{mut}- URA3 [Sn]

5 The fragment PstI-BamHI of the vector Ycplac33-CYR1^{mut}, fragment
 which contains the 3' part of the gene *CYR1*^{mut} (including the *fill* mutation) has been
 sub-cloned in the vector pUC18. Then the auxotrophy marker *URA3*, originating from
 vector pJJ242 (Jones and Prakash (1990) Yeast, 6, pp.363-366) has been inserted in
 this new vector, downstream from the coding part of gene *CYR1*^{mut} in the 3' non
 coding region, at the level of a unique restriction site (*SnaBI*). The obtained vector,
 10 called pUC18-CYR1^{mut}-URA3 is represented in figure 3-3.

b/ Transformation of haploid strains of yeasts M5 and SP1

15 The vector pUC18-CYR1^{mut}-URA3 has been hydrolyzed by the
 appropriate enzyme (here *BalI*), which allowed to obtain a linear fragment of 3.2 kb
 containing the *fill* mutation, the *URA3* marker as well as the sequences located at the
 extremities permitting homologous recombination. This fragment was used for the
 transformation of lab strains : the strain M5 (*ura3*, *trp1*, *leu2*) which is a true haploid
 (n, i.e. 16 chromosomes) originating from the diploid strain M5 (2n) described by
 Schaaf and coll. (1989) Curr. Genet., 15, pp.78-81, and the strain SP1 (*ura3*, *trp1*, *leu2*,
 20 *ade8*) described by Toda and coll. (1985) Cell, 40, pp.27-36.

The presence of the mutation has then been searched for on some of the
 thus obtained transformants by sequencing of the zone carrying in principle the
 mutation. It became thus possible to isolate the strains M5 *fill* and SP1 *fill*.

c/ Search for the phenotype of thermoresistance

25 The thermoresistance of the strains M5 *fill* and SP1 *fill* has been compared
 with that of their respective controls (M5 and SP1) according to the conditions of test
 T1 with an incubation in the presence of glucose for 30 minutes at 30°C and a heat
 treatment of 20 minutes at 52°C.

The results concerning the reconstructed strains M5 *fill* and SP1 *fill*,
 which are collected in table 3-A show that these strains actually present a phenotype of
 30 thermoresistance, which is not observed in the control strains.

SUB
B217

60750-692650

Table 3-A

RESIDUAL VIABILITY AFTER THERMIC SHOCK AT 52°C OF THE
RECONSTRUCTED *fil1* STRAINS AND OF THEIR CONTROLS

Original strain	Residual viability after treatment at 52°C (%)	
	Wild type strain	<i>fil1</i> strain
M5	9 %	96 %
SP1	1 %	16 %

5

d/ Remarks

The *fil1* mutation is a punctual mutation on the *CDC35/CYR1* gene. A deletion of this gene is lethal, the *cyr1* mutants previously described led to very low growth rates.

10

In the mutated *fil1* strains, the mutated gene *CDC35/CYR1* is over-expressed, to compensate the effect of the mutation, considering that the level of AMP cyclic remains low. The response to the *fil1* mutation seems to correspond to a complex metabolite equilibrium, which may depend on the strain genetic background.

15

Furthermore, it must be noted that the *fil1* mutation seems to delay in an important manner the germination of the spores containing this mutation. This deficiency concerning meiose cannot be found in growth or fermentation active phase. It must only be taken into consideration at the level of the constructs of *fil1* strains by classical genetics.

20

EXAMPLE 4 - Characterization of the *fil2* mutation

25

The strain KL1(=W303 *fil2*) has been isolated from the strain W303-1A (Thomas and Rothstein (1989) Cell, 56, pp.619-630) according to a research protocol of *fil* strains similar to those developed in example 1. The *fil2* mutation is monogenic and recessive. In a first step, a thermoresistance test permitted to confirm the improvement obtained by *fil2* mutation. Then a cross-resistance with respect to other stresses has been shown. It was checked that the mutated strain *fil2* was actually a *fil* mutation as that mutated strain was not significantly affected when its growth and its

fermentative ability were compared with those of the original strain W303-1A. Due to the fact that it is a lab strain, the gene carrying the mutation has been searched for.

a/ Resistances to stresses

5 In order to measure the resistance to thermic stresses (heat and freezing), the strains KL1 and W303-1A were cultured on YPD medium at 30°C until obtaining of stationary phase. After centrifugation, the cells were resuspended in YP medium and incubated at 30°C. After 30 minutes, glucose has been added until obtaining a final concentration of 100 mM in the medium. In the case of the heat stress, the
10 incubation at 30°C has been pursued for 30 minutes, and then the cells were incubated at 52°C for 30 minutes. In the case of freezing, the incubation at 30°C has been pursued for 90 minutes, and then the cells were resuspended in ice cold YP medium and frozen at -30°C for 2 days and then thawed at 30°C. Four successive cycles of freezing/thawing have been carried out.

15 These two tests made possible to show that there is a correlation between the thermoresistance and the resistance against freezing in the case of the *fil2* mutation. The strain KL1 has a survival rate of 80% after heat treatment while the survival rate of the control strain W303-1A is of 20 %; furthermore, after freezing in connection with the severe test which has been carried out, the survival rate of the KL1 mutant is
20 almost 4 times higher than that of the control: about 30% for KL1 with respect to about 8% for W303-1A.

b/ Determination of the mutated gene

25 This determination was carried out using a strategy which is identical to that with which it had been possible to identify the *fil1* mutation (cf. example 3). It has thus been determined that the *fil2* mutation was located close to the centromere of chromosome IV. The strain KL1 was consequently complemented with each of the genes located in that region , according to the general strategy disclosed in example 3 :

- transformation of the strain KL1 (auxotrophic for uracil) by centromeric plasmids
30 carrying each one of the genes located at less than 60 kb from the centromere of chromosome X
- search for the transformant clones having lost the phenotype of thermoresistance

- isolation of the gene carrying the mutation starting from mutant KL1 ; retransformation of the strain KL1 with the thus isolated gene in order to verify that it is not a suppressor gene.

The results of the complementation tests have permitted to show that the
 5 *fil2* mutation was located in the *GPR1* gene. This gene codes for a potential receptor coupled to a G protein and it is supposed to initiate the signal pathway associated with G protein encoded by the gene *GPA2* (Xue, Y and coll. (1998) EMBO Journal 17 : 1996-2007).

In order to verify that the phenotype of thermoresistance was actually
 10 associated with a mutation on the gene *GPR1*, the level of thermoresistance of the strain KL1 has been compared with that of the strain W303-1A in which the gene *GPR1* has been deleted (non lethal deletion), as well as with that of the thermosensitive control W303-1A. The results clearly show that the deletion of the *GPR1* gene permits obtaining of a high level of thermoresistance. Furthermore, the diploid obtained by
 15 crossing between KL1 and W303-1A *gpr1Δ* presents a level of thermoresistance as high as that of the strains KL1 and W303-1A *gpr1Δ*, which confirms that the *fil2* mutation actually concerns the *GPR1* gene.

The *GPR1* gene mutated in order to provide the *fil2* phenotype can be isolated according to technique described in example 4 and it permits the
 20 transformation of industrial strains in such a way as to obtain industrial strains having *fil2* phenotype.

In the *fil2* allele of *GPR1* gene, the following mutation point has been found: the base 948 starting from the start codon, that is to say the ATG, is changed, a Thymine (T) becoming an Adenine (A). The codon in the wild type strain is TAT which codes for
 25 a Tyrosine in position 316 in the protein, it is changed in TAA which codes a STOP. The *fil2* allele seems to code for a truncated version of the Gpr1 protein which contains 962 aminoacids. Aminoacid 316 before which the synthesis of the truncated *fil2* protein stops, is located in the 3rd intracellular loop of the receptor protein Gpr1. It is that loop which is known for its interaction in the receptor proteins of mammals with G protein, it is
 30 probably that loop which interacts in the yeast with the Gpa2 protein.

c/ Reintroduction of the mutation in a diploid strain

A true diploid strain, not auxotrophic, relatively close to the industrial strains, has been transformed so that its two genes *GPR1* are replaced by the mutated gene, corresponding to the *fil2* allele. The starting diploid strain and the *fil2* diploid strains have been grown on molasses in fed-batch, in pilot fermentors. A drying test has been performed with the obtained strains as well as a frozen dough stability test of C1 type. The result after drying is a slight decrease of the loss of gassing power with the *fil2* diploid strain compared to the starting strain. A clear improvement by about 1.5 times of the frozen dough stability is shown after 15 days at -20°C.

EXAMPLE 5 - Performances of fil strains during drying

The fil phenotype is interesting for obtaining more active dry yeasts, due to the fact that it permits to subject to drying more active biomasses, without dramatical loss of fermentative ability during drying.

The two reconstructed *fil1* strains, obtained as disclosed in example 3 and the strain S47 *fil500* (= AT28) obtained as described in example 2 and the corresponding starting or original strains have been cultured according to the classical processes in fed batch in pilot fermentor of some liters as described in US patents 4,318,929, 4,318,930 and 4,396,632 in such a way that an active biomass having more than 7% of nitrogen per dry matter is obtained. This biomass is then dried using a process of fluidization type as disclosed in the above-identified US patents.

a/ Drying of the 2 reconstructed *fil1* strains and the 2 original strains M5 and SP1

The fermentative ability of the biomass obtained with each of the 4 strains before drying, i.e. in the form of fresh yeast having about 30% of dry matter has been measured according to test A20. The fermentative ability of the dry yeasts having about 95% of dry matter has been measured according to the test A'20. Due to the fact that, within the frame of such a pilot test, the raw gassing power values do not have any significance, the loss during drying obtained with each original strain has been affected with a coefficient 100 and the loss during drying of the reconstructed *fil1*

strains obtained according to example 3 expressed in percentage of loss during drying of the starting strain. These results are given in table 5A.

Table 5-A

5 EVALUATION OF THE LOSS OF GASSING POWER AFTER DRYING OF THE *fil1* STRAINS
RECONSTRUCTED AND OF THEIR CONTROL

	Loss of ability of the dry yeasts expressed in percentage of the loss of gassing power of the original strain	
	rehydration at 20°C	rehydration at 38°C
M5	100	100
M5 <i>fil1</i>	44	26
SP1	100	100
SP1 <i>fil1</i>	49	52

10 These results show that the *fil1* gene, i.e. the gene *CYR1/CDC35*
carrying the *fil1* mutation allows the construction of industrial strains which present a
loss during drying which is smaller. .

b/ Drying of the strain S47 and of the strain S47 *fil500*

15 The same experiment has been carried out with the strain S47 *fil500*
obtained from a selection based on freeze resistance during the fermentation phase,
taking into account that very often the stress resistances are crossed. The fermentative
abilities have been measured according to test A1 as far as the fresh yeast is concerned,
and according to test A'1 as far as the dry yeast is concerned with rehydration at 20°C.
The results are given in table 5-B.

Table 5-B

EVALUATION OF THE LOSS OF GASSING POWER AFTER DRYING OF THE STRAIN S47 *fil500*
AND OF ITS CONTROL

Loss of gassing power of the dry yeasts expressed in percentage of the loss of gassing power of the starting strain rehydration at 20°C	
S47	100
S47 <i>fil500</i>	66

5

This table 5-B confirms that the fil phenotype allows to diminish the loss during drying of very active yeasts and consequently to obtain very active dry yeasts.

c/ Drying of AT25 = S47 *fil400* and S47 strains

The above experiment has been repeated several times with AT25 = S47 *fil400* strain, which, in all of the tests, proved to be the best strain among AT strains series constructed according to the process described in in example 2.

10

The average results are as follows:

	N/DM Nitrogen on dry matter	A1 test on yeast before drying	A'1 test on active dry yeast at 95% dry matters	Loss due to drying and rehydration at 38°C
S47	8.2%	152	105	31%
AT25	8.3%	165	135	18%

15

EXAMPLE 6 - Use of the fil strains in frozen dough

In this example, are described the results during freezing of fil strains obtained in example 2 : the strain S47 *fil400* (=AT25), the strain S47 *fil500* (=AT28) and the strain AT26. The stability during freezing of these fil strains is compared with that of the strain S47, control strain from which they originate. To do this, the loss of CO₂ production associated with a prolonged storage of frozen pieces of dough at - 20°C has been measured.

20

The yeasts are cultivated on molasses, in fed batch on pilot fermentors in order to obtain active biomasses having a maximum nitrogen content of 8% on dry matter, and then they were used in order to prepare pieces of dough according to the conditions described in the test C1. For each strain, the value of the reference gas (CO₂) production corresponds to the value of the gas production measured after 1 day of conservation at -20°C. The other pieces of dough were thawed after 1 week, 2 weeks, 3 weeks, 4 weeks and 6 weeks ; they allow to monitor the change of the gassing power as a function of the duration of conservation at -20°C. Furthermore, the stability in frozen dough pieces of each strain after n days of conservation at -20°C was measured using the following ratio :

CO₂ release according to C1 test on day n of conservation at -20°C

CO₂ release according to C1 test after the first day of conservation at -20°C.

The results are collected in table 6-A (evolution of the fermentative activity C1 test) and in figure 6-1 (stability in frozen doughs maintained at -20°C).

A very clear improvement of the stability during freezing was noticed for the 3 mutants AT25, AT26 and AT28 (figure 6-1) ; under these conditions of test C1, the control strain S47 maintains after 1 month 54 % of its reference gassing power, while the mutants conserve, under the same conditions between 80 and 87% of their reference gassing power. Furthermore and due to the fact that the mutations did not lead to an important penalization of the fermentative ability, all the mutants attain quickly (from 8 to 15 days of conservation at -20°C) levels of fermentative ability higher than those of control S47 (table 6-A).

sub
B22
15
sub
B23
20

EVOLUTION OF THE GASSING POWER AS A FUNCTION OF THE DURATION OF STORAGE AT -20°C

5

10

In particular, the AT25 strain in test R gives the same result that the S47 strain.

15

20

25

- preservation after 7 days at 26°C in polyethylene plastic bags of at least 40% of the initial activity of freshly produced compressed bakers' yeast in test A1
- no unpleasant smell, no bad taste given to breads, on the contrary, in the processes including a long storage of the frozen doughs (60 days), breads have pure taste with no yeast taste
- better stress resistance to freezing leading in test C1 to a conservation stability defined by the ratio:

$$\frac{\text{CO}_2 \text{ release according to test C1 after a month (30 days) of conservation at } -20^\circ\text{C}}{\text{CO}_2 \text{ release according to test C1 after one day of conservation at } -20^\circ\text{C}} \geq 80\%$$

EXAMPLE 7 - Construction of new fil strains derived from S47 *fil400* = AT25

The genetic analysis of the segregants obtained from AT25 strains led to think that the mutations did concern several genes.

- 15 The following step was then used in an attempt to obtain more performing strains from AT25.

The AT25 segregants were crossed according to their sign with the laboratory haploids W303-1A of mating type a or α . The polyploid strains obtained were subjected to T7 test after 2 weeks of freezing, and classified in this test in comparison with AT25 and S47 and diploid W303-1A.

In this first series of tests, the following values are obtained for the control strains:

	T7 2 weeks	T6 glucose consumption after 90 minutes
AT25	28%	≥ 5.0
S47	14%	≥ 5.0
W303-1A diploid	12%	3.5

27 polyploid strains are obtained with a result equal or superior to S47 strain in T7 test, 7 of which show a result superior to AT25 strain. All these strains have a glucose consumption in T6 test higher than that of diploid W303-1A.

- 25 The segregants issued from AT25, which after crossing with W303-1A haploid revealed a polyploid strain giving in test T7 a result as least equal to S47 strain, are then crossed between them.

Polyploid strains thus obtained are selected with T6 and T7 tests, the selection criteria being as follows:

- glucose consumed within 90 minutes at least equal to 80% of the glucose consumed by AT25 or S47
- 5 - result in T7 after 1 day and after 3 weeks of freezing superior to the result obtained with AT25.

9 polyploid strains corresponding to this definition were obtained, in a first experiment, 3 of which have been deposited at the C.N.C.M. These 3 strains gave the following results:

Strain	T6 glucose consumption in 90 minutes	T7 after 1 day of freezing	T7 after 3 weeks of freezing	cAMP response	T4 μ max. on molasses 0.5%
AT251	5.8	68	38	+	0.6
AT252	5.8	69	40	-	0.6
AT254	5.7	66	40	+	0.6
AT25	5.6	50	26	++	0.6
S47	5.8	28	6	++	0.6

- 10 AT251 strain was deposited at C.N.C.M., 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-....., in accordance with the Budapest Treaty.

AT252 strain was deposited at C.N.C.M., 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-....., in accordance with the Budapest Treaty.

- 15 AT254 strain was deposited at C.N.C.M., 25 rue du Docteur Roux, F-75724 PARIS cedex, under the n° I-....., in accordance with the Budapest Treaty.

These results show that the 3 AT251, AT252 and AT254 strains are good candidates to give in frozen doughs better results than with AT25 strain. These results prove that the process of improvement of an industrial mutant preferably with fil type, and obtained according to the process subject of the invention, bearing mutations on several genes

- 20 compared to the not-mutated industrial strain, consisting:

- in crossing the segregants issued from this industrial mutant with a laboratory haploid strain to select the segregants of the said industrial mutant giving to the polyploids obtained with the laboratory strain an improvement in the required property(ies)

- in crossing the industrial segregants thus selected one with the other and in selecting the polyploids thus obtained according to the criteria of the fil phenotype allows to obtain improved industrial fil strains.

This process is also one of the subjects of the invention, as well as the strains of the
5 same type or kind as the AT251, AT252 and AT254 strains it allows to obtain.

60790 00000000